



## International Journal of Biology Sciences

ISSN Print: 2664-9926  
 ISSN Online: 2664-9934  
 Impact Factor: RJIF 5.45  
 IJBS 2020; 2(2): 68-79  
[www.biologyjournal.net](http://www.biologyjournal.net)  
 Received: 04-08-2020  
 Accepted: 05-09-2020

### Cheng-Bin Wang

Engineering Research Center  
 for Forest and Grassland  
 Disaster Prevention and  
 Reduction, Mianyang Normal  
 University, 166 Mianxing West  
 Road, Mianyang 621000,  
 Sichuan, P. R. China, China

### Tian-Long He

Donghuaxincheng,  
 Wangfenggang, Xiejiaji  
 District, Huainan 232046,  
 Anhui, P. R. China, China

### LI HE

No.66, 5th Shuangcheng Road,  
 Chenghua District, Chengdu  
 610051, Sichuan, P. R. China,  
 China

### Chao Zhou

No. 10, Hengde Road, Jinniu  
 District, Chengdu 610081,  
 Sichuan, P. R. China, China

### Corresponding Author:

#### Cheng-Bin Wang

Engineering Research Center  
 for Forest and Grassland  
 Disaster Prevention and  
 Reduction, Mianyang Normal  
 University, 166 Mianxing West  
 Road, Mianyang 621000,  
 Sichuan, P. R. China, China

## Two new species of *Lucanus* Scopoli, 1763 from Yunnan, Southwest China (Coleoptera, Lucanidae, Lucaninae)

Cheng-Bin Wang, Tian-Long He, LI HE and Chao Zhou

DOI: <https://doi.org/10.33545/26649926.2020.v2.i2a.93>

### Abstract

The stag beetle genus *Lucanus* Scopoli, 1763 (Coleoptera, Lucanidae, Lucaninae, Lucanini) includes 118 species (Without subspecies) worldwide with more than half of them, 64 species, known from China. Members of this genus are very charismatic that they can work their magics on both amateurs and professionals. Two new species of the genus *Lucanus* Scopoli, 1763, *L. niu* sp. n. and *L. zenghuae* sp. n., are described both from Yunnan Province, Southwest China. Diagnostic characters of the two new species are illustrated and comparisons with closely-related congeners are provided in tables.

**Keywords:** stag beetle, lucanini, taxonomy, oriental region

### Introduction

*Lucanus* Scopoli, 1763 is the type genus of Lucanidae (Coleoptera) with members occurring in the Nearctic, Oriental and Palearctic Regions; however, it reaches its highest diversity in eastern Asia. Since the three milestone books of “Stag beetles of China” by Huang & Chen (2010) [5], Huang & Chen (2013) [6] and Huang & Chen (2017) [7, 9], nine species have been contributed to *Lucanus* from Chinese fauna: *L. zhanbishengi* Wang & Zhu, 2017 [12] from Hunan (Wang & Zhu 2017) [12], *L. zhuxiangi* Wang & Zhan, 2018 [13] from Guangdong and Hunan (Wang & Zhan 2018) [13], *L. chengyuani* Wang & Ko, 2018 [14] from Taiwan (Wang & Ko 2018) [14], *L. liuweii* Huang & Chen, 2019 [8] from Zhejiang (Huang & Chen 2019) [8], *L. cenwanglaoshanus* Huang & Chen, 2020 from Guangxi (Huang & Chen 2020), *L. takeoi* Adachi, 2020 from Sichuan (Adachi 2020) [1], *L. yulaoensis* Lin, 2021 from Taiwan (Lin 2021) [10], *L. moae* Qi, 2021 [11] from Sichuan (Qi 2021) [11], and *L. shulini* Bian & Zhan, 2021 [3] from Xizang (Bian & Zhan 2021) [3]. In the present paper, we describe and illustrate two new species of *Lucanus* both from Yunnan Province, Southwest China, under the name of *L. niu* sp. n. and *L. zenghuae* sp. n. Therefore, the number of the *Lucanus* species from China comes to 66 (without subspecies) (Huang & Chen 2010, Huang & Chen 2013 [6], Bartolozzi *et al.* 2016 [5, 2], Huang & Chen 2017 [7, 9], unpublished data of the authors). *L. niu* sp. n. is closely related to *L. vitalisi* Pouillaude, 1913, and *L. zenghuae* sp. n. is most similar to *L. pesarinii* Zilioli, 1998 and *L. derani fukinukiae* Katsura & Giang, 2002. Important morphological characters of the two new species are illustrated and their differential diagnoses from related congeners are provided.

### Materials and Methods

Specimens were relaxed and softened in a digital homeothermic water bath HH-2 at 44.4°C for 12 hours and then placed in distilled water for cleaning, observation and dissection. To examine the genitalia, the abdomens were detached and treated with a 10% potassium hydroxide solution for 12 hours. They were then placed in distilled water to remove the remaining KOH and prevent any further bleaching. After examination, the body parts were mounted on a slide using Euparal Mounting Medium for future studies. Habitus images were taken using a Canon 50D DSLR with a Canon EF 100mm f/2.8L IS USM lens, and a Canon MT-24EX Macro Twin Lite Flash was used as light source. Images of the morphological details were taken with a Canon macro photo lens MP-E 65mm on a Canon 5DsR. Images of the same object at different focal planes were combined using Zerene Stacker 1.04 stacking software. Adobe Photoshop CS6 was used for post-processing.

Morphological terminology follows Holloway (2007) [4], Huang & Chen (2010) [5], Huang & Chen (2013) [6] and Huang & Chen (2017) [7, 9]. The material examined for this study is deposited in the following institutional and private collections: CCZC: collection of Chao Zhou, Chengdu, China; CTLH: Collection of Tian-Long He, Huainan, China; CYCX: collection of Yun-Chuan Xu, Yuxi, China; MAHU: Museum of Anhui University, Hefei, China; MYNU: the invertebrate collection of Mianyang Normal University, Mianyang, China.

The following material was studied for comparison: *Lucanus vitalisi* Pouillaude, 1913 (vs. *Lucanus niu* sp. n.): 3♂♂, CHINA, Yunnan: Puer City, Luchun County, Huanglianshan [普洱市绿春县黄连山], VI.2019, 1800 m, Tian-Long He leg. (CTLH); 4♂♂, same data as previous except: VI.2020; 1♂, VIETNAM: Yen Bai, Mu Cang Chai, V.2017, 1700 m, local people leg. (CTLH); 4♂♂, same data as previous except: V.2019; 3♂♂, same data as previous except: IV.2020. *Lucanus pesarinii* Zilioli, 1998 (vs. *Lucanus zenghuae* Wang & He sp. n.): 1♂, VIETNAM: Vinh Phuc, Tam Dao, VII.2004, local people leg. (CTLH); 1♂, same data as previous except: VI.2021. *Lucanus derani fukinukiae* Katsura & Giang, 2002 (vs. *Lucanus zenghuae* Wang & He sp. n.): 10♂♂, CHINA, Yunnan: Honghe Prefecture, Pingbian County, Daweishan [红河州屏边县大围山], V.2015, Tian-Long He leg. (CTLH); 9♂♂, same data as previous except: V.2016; 12♂♂, same data as previous except: V.2019; 10♂♂, same data as previous except: V.2020; 7♂♂, same data as previous except: V.2021, Tian-Long He & Hao Xu leg.; 3♂♂, CHINA, Yunnan: Honghe Prefecture, Jinping County, Fenshuiling [红河州金平县分水岭], VI.2016, Tian-Long He leg. (CTLH); 2♂♂, same data as previous except: VI.2019; 2♂♂, same data as previous except: VI.2020; 2♂♂, VIETNAM: Lai Chau, Phong Tho, V.2019, local people leg. (CTLH); 4♂♂, same data as previous except: V.2020.

Measurement criteria in millimetres (mm) are as follows: body length: length between the apex of mandible and the elytral apex along the midline; elytral length: length between the basal border and the apex of elytra along suture; elytral width: widest part of both elytra combined; head length: length between the anterior apex of clypeus and the posterior margin of occiput along the midline; head width: widest part of head; mandible length: length from the apex of mandible to its base; pronotal length: length of the pronotum along the midline; pronotal width: widest part of pronotum.

### Taxonomy

*Lucanus niu* Wang, He, He & Zhou, sp. n.

**Material examined.** 11♂♂.

#### Holotype:

♂, CHINA, Yunnan: Puer City, Zhenyuan County [普洱市镇沅县], 12.VI.2020, 2000 m, Shao-You Huang leg. (MAHU).

#### Paratypes

4♂♂, same data as holotype except: (1♂ in CCZC and 3♂♂ in CTLH);

2♂♂, same data as holotype except: 24.VI.2020, Yun-Chuan Xu leg. (1♂ in CTLH and 1♂ in CYCX);

4♂♂, same data as holotype except: 5.VI.2021, 2000 m, local people leg. (3♂♂ in MAHU and 1♂ in MYNU).

**Description.** Holotype male. Large size, body 57.4 mm long. Lengths of body parts (mm): head (9.3), mandible (21.2), pronotum (8.3), elytra (23.0); width (mm): head (14.8), pronotum (13.6), elytra (16.6).

**Habitus** (Fig. 1a–c). Color mostly reddish brown to blackish brown; elytra reddish brown; femora with reddish stripes on ventral surfaces; tibiae with obscure reddish stripes on ventral surfaces; protibiae tinged with reddish on dorsal surfaces. Body mostly clothed with short, sparse, recumbent, yellowish pubescence, similar on metaventre, much finer and denser on elytra.

Head 1.6 times as wide as long, widest at anterolateral angles; surfaces matt, finely punctate, intervals finely and irregularly granulose. Frons relatively long, fan-like, weakly concave, fused with clypeolabrum. Clypeal ridge strong, medially interrupted, laterally projecting as two ridges. Labrum slenderly subtriangular, rounded at apex. Anterolateral angles acute, strongly protruding laterally, more prominent than eyes. Ocular canthus protruding, shorter than half eye's diameter. Postocular margin almost oblique, gradually narrowed posteriorly. Posterior margin widely concave. Anterior ridge distinctly and oblongly elevated, substraight at apical margin. Lateral ridges rather strongly elevated, broad, widely subrounded at apical margins. Mandible long and slender, about 2.3 times as long as head; in dorsal view, mandible curved in basal 1/4 and apical 1/5, and almost straight in between; in lateral view, mandible strongly curved in basal 3/5 and almost straight in apical 2/5; surfaces finely punctate, intervals matt and finely granulose at inner side, feebly sheen and microreticulate at outer side. Inner mandibular margin with unarmed carina at base, preceded by diastema, 7–8 small denticles, major inner tooth, and 2–3 distal small denticles. Major inner tooth situated at apical 1/3, strong, digitiform, shorter than mandibular width, tilting forwards and upwards with apex rounded and slightly higher than mandible. Apical fork with upper tooth bent upwards and inwards; lower tooth markedly shorter, conical, slightly directed downwards and inwards. Antennal club 4-segmented; antennomere VII with much thinner lamella, sharply pointing inwards, only pubescent on apical surface; antennomeres VIII–X with strong, long, pubescent lamellae. Mentum wide, inverted trapezoidal; anterior margin almost straight and with obliquely truncated corners; surface matt, finely punctate, intervals finely and irregularly granulose, and transversely depressed near anterior margin.

Pronotum 1.6 times as wide as long, widest at middle. Anterior corners weakly acute; posterior corners obtuse. Surface feebly sheen, finely punctate, intervals finely and irregularly granulose, and with median longitudinal depression rather shallow. Prosternal apophysis strong, distinctly convex ventrally in middle of posterior part, and truncated at posterior margin.

Scutellum linguiform, about 1.8 times as wide as long, rounded at apical margin. Surface feebly sheen, finely punctate, intervals microreticulate.

Elytra oval, 1.4 times as long as wide, widest around basal 4/9, 2.8 times as long as and 1.2 times as wide as pronotum. Humeri rounded. Surface sheen, finely and densely punctate, intervals microreticulate.

Legs slender. Protibia with 5 (left, basal one tiny) or 4 (right, basal one tiny) teeth along outer margin; apex

bifurcate with branches narrowly rounded at tips. Except apical spurs and spines, mesotibia with 3 (right) or 4 (left, basal one tiny) small lateral spines and metatibia with 3 lateral spines (basal one tiny). Surfaces finely punctate, intervals matt and finely granulose on femora and tibiae, feebly sheen and microreticulate on tarsi.

Male genitalia. Abdominal tergite VIII (Fig. 2a) with subround lateral angles and with longitudinal, thin, membranous stripe in middle of basal 5/9; sternite VIII (Fig. 2b) with transverse, semielliptical, membranous area in middle of basal half. Abdominal sternite IX (Fig. 2c) with apical expansion relatively slender, and with long, membranous, median stripe in apical 5/8; basal stalk not extended laterally. Aedeagus (Fig. 2f) about 2.3 times as long as wide in ventral view. Basal piece (Fig. 2f–h) constricted in basal part, about 1.8 times as long as parameres, with a pair of sclerotized dorsal plates (Fig. 2h); ventral plate (Fig. 2g) bifurcate at apex of distal end, with relatively deep emargination. Paramere with basal process (Fig. 2g–h); apex strongly upturned in lateral view (Fig. 2i). Penis (Fig. 2f–h) moderately wide, weakly expanded near apex, 0.2 times shorter than parameres. Flagellum relatively short, about 2.2 times as long as parameres, apex distinctly enlarged.

Male paratypes. Body 43.5–58.0 mm long. All male paratypes consistent with holotype except for usual allometric differences (Fig. 1d–i). Large-sized male like holotype with more pronounced morphological characters, while in small-sized male, mandible strongly shorter, apical fork of mandible less opened to vestigial, number of inner teeth fewer, major inner tooth ill-developed, clypeolabrum shorter, anterior and lateral ridges of head weaker.

Female. Unkown.

**Etymology.** The specific epithet is from the Chinese Pinyin “niú [牛]”, which means ox, indicating this species was identified as new in 2021, the year of the ox. The name is a noun in apposition.

**Distribution.** China (Yunnan).

**Differential diagnosis.** This new species should be assigned to the *Lucanus laminifer* group (*sensu* Huang & Chen 2010)<sup>[5]</sup>. It most resembles *Lucanus vitalisi* Pouillaude, 1913, but can be distinguished by the combination of characters shown in Table 1.

### ***Lucanus zenghuae* Wang, He, He & Zhou, sp. n.**

**Material examined.** 6♂♂.

#### **Holotype:**

♂, CHINA, Yunnan: Honghe Prefecture, Pingbian County, Daweishan [红河州屏边县大围山], 5.VI.2020, Tian-Long He leg. (MAHU).

#### **Paratypes**

2♂♂, same data as holotype except: 9.VI.2019 (CTLH);

2♂♂, same data as holotype except: 31.V.2021, Tian-Long He & Hao Xu legg. (1♂ in CTLH and 1♂ in MYNU);

1♂, same data as holotype except: 31.V.2021, Tian-Long He & Hao Xu legg. (CCZC).

**Description.** Holotype male. Large size, body 48.6 mm long. Lengths of body parts (mm): head (8.4), mandible (15.9), pronotum (6.4), elytra (19.7); width (mm): head (15.2), pronotum (11.0), elytra (13.2).

**Habitus** (Fig. 5a–c). Color mostly reddish brown to blackish brown; elytra reddish; femora and tibiae with reddish stripes

on ventral surfaces. Body mostly clothed with short, sparse, recumbent, yellowish pubescence, similar on metaventrite.

Head 1.8 times as wide as long, widest at anterolateral angles; surfaces matt, finely punctate, intervals finely and irregularly granulose. Frons transverse, subtrapezoidal, weakly concave, fused with clypeolabrum. Clypeal ridge strong, not medially interrupted, laterally projecting as two teeth. Labrum subtriangular, rounded at apex. Anterolateral angles subacute, protruding laterally, slightly prominent than eyes. Ocular canthus protruding, shorter than half eye's diameter. Postocular margin gently curved, strongly narrowed posteriorly. Posterior margin widely concave. Anterior ridge weakly elevated, transverse, substraight at apical margin. Lateral ridges moderately elevated, broad, widely subrounded at apical margins. Mandible moderately long and strong, about 1.9 times as long as head; moderately curved in dorsal and lateral views; surfaces finely punctate, intervals matt and finely granulose at inner side, matt and microrugulate at outer side. Inner mandibular margin with unarmed carina at base, preceded by 5–6 small denticles, major inner tooth, and 4–5 distal small denticles. Major inner tooth situated at apical 2/5, rather strong, slenderly subtriangular, weakly protruded near base of posterior side, shorter than mandibular width, tilting backwards and upwards with apex rounded and not higher than mandible. Apical fork with upper tooth bent upwards and inwards; lower tooth distinctly longer, directed downwards and inwards. Antennal club 4-segmented; antennomere VII with slenderly subtriangular lamella, sharply pointing inwards, only pubescent on apical surface; antennomeres VIII–X with strong, long, pubescent lamellae. Mentum wide, inverted trapezoidal; anterior margin almost straight and with obliquely truncated corners; surface matt, finely punctate, intervals finely and irregularly granulose, and transversely depressed near anterior margin.

Pronotum 1.7 times as wide as long, widest at middle. Anterior corners subrounded; posterior corners obtuse. Surface matt, finely punctate, intervals finely and irregularly granulose, and with median longitudinal depression rather shallow. Prosternal apophysis strong, distinctly convex ventrally in middle of posterior part, and truncated at posterior margin.

Scutellum linguiform, about 1.9 times as wide as long, rounded at apical margin. Surface feebly sheen, finely punctate, intervals microrugulate.

Elytra oval, 1.5 times as long as wide, widest around basal 1/3, 3.1 times as long as and 1.2 times as wide as pronotum. Humeri obtuse. Surface weakly sheen, finely and densely punctate, intervals microreticulate.

Legs slender. Protibia with 3 (left, basal one tiny) or 4 (right, basal two tiny) teeth along outer margin; apex bifurcate with branches narrowly rounded at tips. Except apical spurs and spines, mesotibia with 3 (basal one tiny) small lateral spines and metatibia without lateral spines (only one unobscured protuberance). Surfaces finely punctate, intervals matt and finely granulose on femora and tibiae, feebly sheen and microreticulate on tarsi.

Male genitalia. Abdominal tergite VIII (Fig. 6a) without lateral angles and with longitudinal, thin, membranous stripe in middle of basal 5/8; sternite VIII (Fig. 6b) with transverse, semielliptical, membranous area in middle of basal part. Abdominal sternite IX (Fig. 6c, e) with apical expansion relatively slender, without membranous area; basal stalk not extended laterally. Aedeagus (Fig. 6f) about

3.0 times as long as wide in ventral view. Basal piece (Fig. 6f, g) weakly constricted in basal part, about 1.4 times as long as parameres, with a pair of sclerotized dorsal plates (Fig. 6g); ventral plate (Fig. 6f) bifurcate at apex of distal end, with relatively deep emargination. Paramere with basal process (Fig. 6g); apex moderately upturned in lateral view (Fig. 6h). Penis (Fig. 6f) slender, 0.4 times shorter than parameres. Flagellum relatively short, about 2.2 times as long as parameres, apex distinctly enlarged.

Male paratypes. Body 34.5–49.1 mm long. All male paratypes consistent with holotype except for usual allometric differences. Large-sized male like holotype with more pronounced morphological characters, while in small-sized male, mandible shorter, apical fork of mandible less opened, number of inner teeth fewer, major inner tooth ill-developed, clypeolabrum shorter, anterior and lateral ridges of head weaker.

Female. Unkown.

Etymology. The new species is dedicated to Zeng-Hua Tang, the mother of the collector Tian-Long He. The name is a noun in the genitive case.

Distribution. China (Yunnan).

Differential diagnosis. This new species should be assigned to the *Lucanus fortunei* group (*sensu* Huang & Chen 2010)


[5]. It is most similar to *L. pesarinii* Zilioli, 1998 and *L. derani fukinukiae* Katsura & Giang, 2002, but can be distinguished by the combination of characters shown in Table 2.

#### Acknowledgements


We appreciate Gui-Qiang Huang (Liupanshui Normal University, Liupanshui, China), who rekindling our enthusiasm for taxonomy. We would like to express our sincere gratitude to Chang-Chin Chen (Tianjin, China), Hao Huang (Qingdao, China), Mei-Ying Lin, Yun-Long Ma, Lu Qiu, Jian-Yue Qiu and Hao Xu (MYNU), Liang Lü (Hebei Normal University, Shijiazhuang, China), Xia Wan (Anhui University, Hefei, China), Ping Wang and Guang-Lin Xie (Yangtze University, Jingzhou, China), Yun-Chuan Xu (Yuxi, China), Jian-Bo Yang (Dali, China), Qiao-Zhi Yang (University College London, U.K.), Mao Ye (Xiangyang, China), Bi-Sheng Zhan (Zhenjiang, China), Zheng Zhou (Sichuan Agricultural University, Chengdu, China), and Xiang Zhu (Beijing, China) for their considerable help in our study. We are grateful to reviewers who provided constructive comments on an earlier version of the manuscript. This study was supported by the Natural Science Foundation of Huaguoshan (NSFH-2021).




**Fig 1:** Male habitus of *Lucanus niu* sp. n. a–c large size, holotype; d–f moderate size, paratype; g–i small size, paratype. a, d, g: dorsal view; b, e, h: ventral view; c, f, i: lateral view. Scale bar = 1.0 cm.




**Fig 2:** Male genitalia of *Lucanus niu* sp. n., holotype. a abdominal tergite VIII; b abdominal sternite VIII; c–e abdominal segment IX; f–i aedeagus (penis removed in g). a, d, h: dorsal view; b, c, f, g: ventral view; e, i: lateral view. Scale bar = 1.0 mm.




**Fig 3:** Male habitus of *Lucanus vitalisi* Pouillaude, 1913 from Yunnan, China. a–c large size; d–f moderate size; g–i small size. a, d, g: dorsal view; b, e, h: ventral view; c, f, i: lateral view. Scale bar = 1.0 cm.




**Fig 4:** Male genitalia of *Lucanus vitalisi* Pouillaude, 1913 from Yunnan, China. a abdominal tergite VIII; b abdominal sternite VIII; c–e abdominal segment IX; f–i aedeagus (penis removed in g). a, d, h: dorsal view; b, c, f, g: ventral view; e, i: lateral view. Scale bar = 1.0 mm.




**Fig 5:** Male habitus of *Lucanus zenghuae* sp. n., holotype. a dorsal view; b ventral view; c lateral view. Scale bar = 1.0 cm.




**Fig 6:** Male genitalia of *Lucanus zenghuae* sp. n., holotype. a abdominal tergite VIII; b abdominal sternite VIII; c-e abdominal segment IX; f-h aedeagus. a, d, g: dorsal view; b, c, f: ventral view; e, h: lateral view. Scale bar = 1.0 mm.




**Fig 7:** Male habitus of *Lucanus pesarinii* Zilioli, 1998 from Vietnam. a dorsal view; b ventral view; c lateral view. Scale bar = 1.0 cm.




**Fig 8:** Male genitalia of *Lucanus pesarinii* Zilioli, 1998 from Vietnam. a abdominal tergite VIII; b abdominal sternite VIII; c–e abdominal segment IX; f–h aedeagus. a, d, g: dorsal view; b, c, f: ventral view; e, h: lateral view. Scale bar = 1.0 mm.





**Fig 9:** Male habitus of *Lucanus derani fukinukiae* Katsura & Giang, 2002 from Yunnan, China. a dorsal view; b ventral view; c lateral view. Scale bar = 1.0 cm.



**Fig 10:** Male genitalia of *Lucanus derani fukinukiae* Katsura & Giang, 2002 from Yunnan, China. a:abdominal tergite VIII; b abdominal sternite VIII; c–e abdominal segment IX; f–h aedeagus. a, d, g: dorsal view; b, c, f: ventral view; e, h: lateral view. Scale bar = 1.0 mm.

**Table 1:** Selected important morphological differences between males of *Lucanus niu* sp. n. and *L. vitalisi* Pouillaude, 1913.

		<i>Lucanus niu</i> sp. n.	<i>Lucanus vitalisi</i> Pouillaude, 1913
Body length		large-sized male shorter than 60 mm (Fig. 1a–c)	large-sized male generally longer than 70 mm (Fig. 3a–c)
Color	femora	reddish stripes distinct on ventral surfaces in large-sized male (Fig. 1b), but obscure in moderate-sized and small-sized males (Fig. 1e, h)	reddish stripes distinct on ventral surfaces from large-sized to small-sized males (Fig. 3b, e, h)
	tibiae	reddish stripes obscure on ventral surfaces (Fig. 1b, e, h), and absent on dorsal surfaces (protibiae tinged with reddish) (Fig. 1a, d, j)	reddish stripes distinct on ventral surfaces (Fig. 3b, e, h), and obscure on dorsal surfaces (Fig. 3a, d, j)
Head	mandibles	about 2.3 times as long as head in large-sized male (Fig. 1a–c); inner mandibular margin with unarmed carina at base, preceded by diastema, 7–8 small denticles, major inner tooth, and 2–3 distal small denticles in large-sized male (Fig. 1a, b); major inner tooth relatively more prominent in large-sized male (Fig. 1a, b); apical part thick and strong in moderate-sized and small-sized males (Fig. 1d, e, g, h)	about 3.0 times as long as head in large-sized male (Fig. 3a–c); inner mandibular margin with unarmed carina at base, preceded by diastema, 17–18 small denticles, major inner tooth, and 7–8 distal small denticles in large-sized male (Fig. 3a, b); major inner tooth relatively less prominent in large-sized male (Fig. 3a, b); apical part much more slender in moderate-sized and small-sized males (Fig. 3d, e, g, h)
	anterior ridge	narrower; substraight at apical margin	broader; rounded at apical margin
Male genitalia	abdominal sternite VIII	membranous area semielliptical (Fig. 2b)	membranous area trapezoidal (Fig. 4b)
	abdominal sternite IX	apical expansion relatively slender, and with long, membranous, median stripe in apical 5/8 (Fig. 2c)	apical expansion relatively broad, and with short, membranous, median stripe in apical 1/3 (Fig. 4c)
	basal piece	apex of distal end with relatively deep emargination (Fig. 2g)	apex of distal end with relatively shallow emargination (Fig. 4g)
	parameres	apex strongly upturned in lateral view (Fig. 2i)	with apex less strongly upturned in lateral view (Fig. 4i)
	penis	moderately wide, weakly expanded near apex (Fig. 2f)	rather wide, strongly expanded near apex (Fig. 4f)

**Table 2:** Selected important morphological differences among males of *Lucanus zenghuae* sp. n., *L. pesarinii* Zilioli, 1998 and *L. derani fukinukiae* Katsura & Giang, 2002.

		<i>Lucanus zenghuae</i> sp. n.	<i>Lucanus pesarinii</i> Zilioli, 1998	<i>Lucanus derani fukinukiae</i> Katsura & Giang, 2002
Color	elytra	reddish (Fig. 5a, c)	orange-red (Fig. 7a, c)	reddish brown to blackish brown (Fig. 9a, c)
	femora & tibiae	reddish stripes distinct on ventral surfaces (Fig. 5b, c)	orange-red stripes distinct on ventral surfaces (Fig. 7b, c)	reddish stripes absent or vestigial on ventral surfaces (Fig. 9b, c)
Pubescence	metaventricle	short and sparse as on other parts (Fig. 5b)	short and sparse as on other parts (Fig. 7b)	longer and denser than on other parts (Fig. 9b)
Head	major inner tooth	tilting backwards and upwards with apex not higher than mandible (Cannot be seen in lateral view) (Fig. 5a, c)	tilting forwards and upwards with apex distinctly higher than mandible (can be seen in lateral view) (Fig. 7a, c)	tilting backwards and upwards with apex distinctly higher than mandible (can be seen in lateral view) (Fig. 9a, c)
	distal small denticles	continuous with major inner tooth (Fig. 5a)	distinctly spaced out with major inner tooth (Fig. 7a)	distinctly spaced out with major inner tooth (Fig. 9a)
	apical fork	upper tooth more bent inwards, shorter than lower tooth (Fig. 5a)	upper tooth less bent inwards, shorter than lower tooth (Fig. 7a)	upper tooth more bent inwards, not shorter than lower tooth (Fig. 9a)
	anterior ridge	weakly elevated, transverse, substraight at apical margin	absent	weakly elevated, transverse, gently curved at apical margin
Legs	protibia	apical branches less open (Fig. 5a)	apical branches more open (Fig. 7a)	apical branches more open (Fig. 9a)
Male genitalia	abdominal sternite VIII	membranous area transverse, semielliptical (Fig. 6b)	membranous area transverse, semielliptical (Fig. 8b)	membranous areas bisected into two small parts (Fig. 10b)
	abdominal sternite IX	basal stalk not extended laterally (Fig. 6c, d)	basal stalk extended laterally (Fig. 8c, d)	basal stalk extended laterally (Fig. 10c, d)
	penis	slender, not expanded near apex (Fig. 6f)	wide, expanded near apex (Fig. 8f)	wide, expanded near apex (Fig. 10f)

## References

- Adachi N. A new species of the genus *Lucanus* from Szechuan province, southwest China. *Kogane*. 2020;23:8-10.
- Bartolozzi L, Sprecher-Uebersax E, Bezděk A. Family Lucanidae Latreille, 1804. In: Löbl, I. & Löbl, D. (Eds.), *Catalogue of Palaearctic Coleoptera. Scarabaeoidea-Scirtoidea-Dascilloidea-Buprestoidea-Byrrhoidea*. Revised and Updated Edition. Brill, Leiden and Boston. 2016;3:58–84. [https://doi.org/10.1163/9789004309142\\_003](https://doi.org/10.1163/9789004309142_003)
- Bian CZ, Zhan ZH. A new species of *Lucanus* Scopoli, 1763 (Coleoptera, Lucanidae) from Tibet, China. *Fauniversity*. 2021;9(27):1-4.
- Holloway BA. Lucanidae (Insecta: Coleoptera). *Fauna of New Zealand*. 2007;61:1-254.
- Huang H, Chen CC. Stag beetles of China I. Formosa Ecological Company, Taipei, 2010, 288.
- Huang H, Chen CC. Stag beetles of China II. Formosa Ecological Company, Taipei, 2013, 716.
- Huang H, Chen CC. Stag beetles of China III. Formosa Ecological Company, Taipei, 2017, 524.

8. Huang H, Chen CC. *Lucanus liuweii* sp. nov. (Coleoptera: Lucanidae: Lucaninae) from Zhejiang, China. *Beetles World*. 2019;19:2-6.
9. Huang H, Chen CC. New or little-known stag beetles (Coleoptera: Lucanidae: Lucaninae) from China. *Beetles World*. 2017;21:2-11.
10. Lin JZ. Description of *Lucanus yulaoensis* sp. nov., a new species stag beetle from northern Taiwan (Coleoptera, Lucanidae). *Faunitaxys*. 2021;9(2):1-5.
11. Qi ZH. *Lucanus moae* sp. nov., A new species from Sichuan, China (Coleoptera: Lucanidae: Lucaninae). *Faunitaxys*. 2021;9(23):1-7.
12. Wang CB, Zhu X. *Lucanus zhanbishengi* sp. n., a new species from Hunan, central China (Coleoptera: Lucanidae: Lucaninae). *International Journal of Research Studies in Zoology*. 2017;3(4):54-69. <http://dx.doi.org/10.20431/2454-941X.0304008>
13. Wang CB, Zhan BS. *Lucanus zhuxiangi* sp. n., a new species from southeast China (Coleoptera: Lucanidae: Lucaninae). *International Journal of Research Studies in Zoology*. 2018;4(3):1-11. <https://doi.org/10.20431/2454-941X.0403001>
14. Wang LJ, Ko HP. Description of *Lucanus chengyuani* sp. nov. from Taiwan, with a key to the species of Taiwanese *Lucanus* Scopoli (Coleoptera: Lucanidae). *Japanese Journal of Systematic Entomology*. 2018;24(2):257-263.