



## The assessment of floral abundance and composition of Neni-Nimo watershed in Anaocha L.G.A. of Anambra state, Nigeria

Ezenwata Ifeoma Susan<sup>1</sup>, Okemadu Obioma Christian<sup>1</sup>, Nweze Kenneth Emeka<sup>2</sup>, Eze Chinwe Catherine<sup>2</sup>, Onyemeka Regland Michael<sup>3</sup>

<sup>1</sup> Department of Biological Sciences, Chukwuemeka Odumegwu Ojukwu University, Anambra, Nigeria

<sup>2</sup> Department of Microbiology, Federal University of Technology, Owerri, Nigeria

<sup>3</sup> Department of Botany, Lagos State University, Lagos, Nigeria

DOI: <https://doi.org/10.33545/26649926.2021.v3.i1a.20>

### Abstract

The research was carried out on the floral composition of Neni-Nimo watershed between November 2019 and July 2020. The research work aimed to find out the various plant species that inhabit the watershed, their abundance measures and importance values. In this study, the watershed was divided into three catchment areas, the upper, the middle and the lower catchment. Each of the catchment zones was divided into component parts, with an area of (50 m X 30 m) each. The experiment was laid in a randomized complete block design. A plotless sample technique called the centre point technique was used to ascertain importance values Indices (IVI) for trees, while the plot count sampling technique (quadrant) was employed for shrubs, forbs, grasses and climbers. Random sampling techniques were employed in all cases to eliminate bias. The encountered growth forms such as trees, shrubs, forbs, climbers and grasses were identified and recorded. The important values index (IVI) was obtained for all the growth forms. The encountered species were quantitatively analyzed for density, frequency and abundance or dominance depending on their various growth forms. The importance value indices for each of the species were determined by summing up relative density, relative frequency and relative abundance of the species. The result shows that trees were the most abundant flora in the upper catchment area with *Elaeis guineensis* being the most dominant species. At the middle catchment, shrubs were the most abundant flora, with *Chromolaena odorata* being the most dominant species. Also at the lower catchment area, shrubs were the most abundant flora, followed by grasses. *Venonia amygdalina* had the highest abundance for shrubs while *Zea mays* had the most abundance value for grasses at the lower catchment. The study revealed that most of the dominant species in the three catchment areas were crops of economic value which are planted by man.

**Keywords:** catchment, abundance measures, and importance value

### Introduction

A watershed is a topographically delineated area that is bounded by a drainage divide, which collects run-off from precipitation and primarily drains large amounts of underground water, melted ice, nutrients, sediments and toxins, from the soil into the nearby water bodies or stream channels. The term watershed is synonymously used with a drainage basin although they are different (Enwelu *et al.*, 2014) [6]. Watershed is made up of a riparian area or riparian zone which is the strip of vegetation along a stream with distinct composition and density from the surrounding upland. Because of its topographic nature, it drains underground water and nutrients into nearby streams from the predominant landscape surface types (Kunstler *et al.*, 2004) [10]. The impacts of anthropogenic activities and nutrient fluxes from watersheds are dynamic over time and space. Nutrients from watersheds will result in to increase in nutrient in the nearby or receiving water (Howarth *et al.*, 2012; Boyer *et al.*, 2002.) [9, 3]. This nutrient enrichment vary to time and space, depending on many social, regulatory, economic and environmental drivers (Boyer *et al.*, 2002; Russel *et al.*, 2008;

Hale *et al.*, 2013) [3, 14, 8] and also determines the type of flora in that watershed. Most watersheds are usually thick and evergreen forest habitat, with varieties of flora and fauna composition.

The activities of man have led to alterations of the environment which are both passive and diverse. These alterations range from altered atmospheric chemistry to dramatic modifications of the landscape for agricultural and urban uses (Vitousek *et al.*, 1997; Kareiver *et al.*, 2007 [15]. Increasingly, it has been recognized that human actions on watersheds such as deforestation and agricultural activities have created impacts on the ecosystem and is highly connected to the enrichment of nutrient in water bodies (Foley, 2005) [7].

Assessment of species composition entails studies of different plant life as they are naturally distributed on the watershed and their importance value. Importance value is the measure of how dominant a species is in a given forest area. It is usually calculated by summing up the relative frequency, relative density and relative dominance for trees and relative frequency, relative density and relative abundance for shrubs,

forbs and grasses. Bhadra and Pattanayak (2017) [2] opined that dominance is justified for tree community as the individuals of tree species are available in higher girth and abundance is justified for shrubs, forbs and trees. Therefore important value index of species belonging to a tree and other communities should be evaluated based on dominance and abundance separately. Abundance and dominance are two aspects of plant species dynamic, often considered interchangeably together with frequency and density to estimate important value index (IVI) Importance value (IVI). Studies and knowledge of the specie composition of any area especially watersheds are important as it helps to encourage conservation of species. Conservation of biological diversity is essential for the survival of the human race, because it provides a livelihood for man and livestock, mitigate climatic changes and maintain samples of unchanged biotic communities in their natural form for breeding. Therefore, this research work aimed to find out the various plant species that inhabit the watershed, their abundance measures and importance values.

## Materials and Method

### Experimental site

The study on species compositions was carried out at Ogbujilekwe stream watershed bordering Neni and Nimo town in Anaocha L.G. A. of Anambra State, Nigeria. The Ogbujilekwe stream, as explained by the villagers and some of the farmers in the area, cuts across four major towns in Anambra State namely; Neni, Nimo, Oraukwu and Adazi-Nnukwu. The Neni-Nimo watershed falls within the humid tropical climate belt of Nigeria. The two seasons experienced in this area are the rainy season and dry season. The maximum average rainfall is experienced during July and August. The mean annual rainfall is about 1500 – 2500mm.


## Experimental Design

The experiment was laid in a randomized complete block design (RCBD). The watershed was divided into three catchment areas, the upper, middle, and lower catchment, each of the catchments, had three replicates with an equal area of 50m x 30m. The catchments were demarcated from each other, using pegs. The catchment zones were labelled as sites A, B and C. Site A is the upper course/catchment, site B is the middle course and C is the lower catchment. AR1 denote catchment A replicate1, AR2 denote catchment A replicate 2, AR3 denote catchment A replicate 3, BR1 denote catchment B replicate 1, BR2 denote catchment B replicate 2, BR3 denote catchment B replicate 3, CR1 denote catchment C replicate 1, CR2 catchment C replicate 2, CR3 denote catchment C replicate 3, AR4 denote catchment A control sample, BR4 denote catchment B Control sample while CR4 catchment C control sample.

## Sampling procedure

A quantitative relationship between area and number of species was first developed by Arrhenius (1921) [1], a pioneer in the area of Geographical ecology. Point centred plotless count technique was used for the sampling trees, while the plot count technique using quadrants were used for the forbs, shrubs, climbers and grasses. The species encountered were identified using Google lens applicaion, relevant texts like Handbook of West African Weed (Akobundu and Agyakwa, 1998). The vegetation data collected were quantitatively analyzed for density, frequency, relative density, relative frequency and basal area as per (Mishra, 1968) [11]. The importance value index (IVI) for each species was determined as the sum of relative density, relative frequency and relative dominance for the forested area or relative abundance for the non-forested areas (Phillips, 1959).

## Survey Procedure for Trees



**Fig 1:** Diagram representation of the point centred plotless count technique for easier appreciation. Plant species found at numbers 1, 2, 3 and 4 are the closest individual from the sampling points.

### Details of Standard Procedure

The plotless sampling procedure was used for data collection for all trees. Four quarters were established at the sampling point using a cross made by two wooden frames. Firstly, the distance of the closest tree to the sampling point from four quadrants were measured. The averages of the four distances were calculated to obtain the mean distance occupied by trees. Finally, any tree whose stem was up to 4.3ft height had the girth measured immediately at this mark. The reliability of this method has been given by Cottan and Curtix (1994). Okereke *et al.* (2016) <sup>[12]</sup> noted that one of the limitations of this method for field application is that an individual must be located within each quarter and an individual must not be measured twice. The diameter of a tree at 4.3 foot level is called diameter breast height (DBH). The basal area of the trees were calculated using the formulae  $BA = 0.005454 \times DBH^2$  and it was used to ascertain the basal area of the plants. The number 0.005454 is called foresters constant and it comes by dividing  $\pi$  by  $4 \times 144$ . The number of individuals of each species were added up and used to estimate species composition and diversity. Bar charts were used to record species composition according to growth forms (tree, climbers, shrubs, grasses and forbs

### Details of Standard Procedure for Shrubs, Forbs and Climbers

The plot count technique was achieved using the quadrant. The Plot count quadrant techniques are used in herbaceous sites with known boundaries and lacking physical obstacles. At first, the total length and width are obtained. The plots were then demarcated from each other using pegs and the size was ascertained. Then the sampling intensity was worked out by determining 5 % of the total plot size since ecology accepts of 5 % or 10 % for expediency. The sampling unit is a quadrant. Having known the sampling intensity, the number of quadrants to be placed in each of the plots was calculated by dividing sampling intensity by the size of the quadrant. The quadrant size in all cases is 2x1m in size.

Next is to determine the sampling points on the plot. This was achieved by using the random number technique. In this case, a set of random numbers were written according to the number of throws of the quadrant which is 38 the numbers were randomly picked before the experiment and the point of intersection of the numbers were marked with pegs.

### The formulae for calculating the important value indices (IVI) for trees is as follows

$$\text{Density} = \frac{\text{Total number of individual species}}{\text{Total area of the plot sampled}} \quad (1.1)$$

$$\text{Relative density of each species} = \frac{\text{No of the species}}{\text{No of all trees}} \times \text{density of all trees}$$

$$\text{Frequency} = \frac{\text{Number of sampling point at which each specie is found}}{\text{Total no of sampling point}} \times \frac{100}{1}$$

$$\text{Relative frequency} = \frac{\text{Frequency of one spp}}{\text{Total frequency of all species}} \times \frac{100}{1}$$

Dominance of each specie = its relative density  $\times$  its basal area

$$\text{Basal area (BA)} = 0.005454 \times \text{DBH}^2$$

$$\text{Relative dominance} = \frac{\text{Dominance of each specie}}{\text{Total dominance of all species!}} \times \frac{100}{1} \quad (1.2)$$

Importance value IVI = Rel. density + rel. frequency + rel. dominance (COX, 1976).

Formulae for calculation of important value indices IVI of shrubs, trees, and climbers is.

$$\text{Density} = \frac{\text{Total number of individual species}}{\text{Total area of the plot sampled}} \quad (1.3)$$

$$\text{Frequency} = \frac{\text{Number of times each spp occurred}}{\text{Total area of the plots sampled}} \times \frac{100}{1}$$

$$\text{Relative density} = \frac{\text{density of one spp}}{\text{Total density of all species}} \times \frac{100}{1}$$

$$\text{Relative density} = \frac{\text{Frequency of one spp}}{\text{Total frequency of all species}} \times \frac{100}{1}$$

Importance value indices (IVI) = Relative density of species + Relative frequency + Relative abundance

Where:

Density is defined as the number of individuals of particular species per unit area. It also gives an idea of how closely packed trees are growing in an area. Frequency is defined as the distribution or relative representation of a particular species or how species are spread out in a given area. Dominance is defined as the measure of the size or weight of vegetation. Abundance is defined as the number of individual species found in a particular sample.

Basal area is defined as the sum of the cross sectional area of all the trees of a particular species measured at 4.3 ft above the ground. DBH is diameter at breast height of a tree species at 4.3 ft, and 0.005454 is known as foresters constant.

### Result and Discussion

#### Species composition of the studied Watershed

The specie composition of the plants encountered in the three catchment zones was evaluated. A total of 149 plant species comprising of 40 trees, 30 shrubs, 39 forbs, 27 grasses and 13 forbs were encountered. The plant species encountered make up to 45 families. All the growth forms which include trees, forbs, shrubs, climbers and grasses were present in the upper catchment areas or forested area, and middle catchment or fallow areas, but in the lower catchment or current usage area, trees were totally absent.

Looking at Table 1 below, Out of the 40 trees encountered in the watershed, *Elias guinensis*, had the highest dominance value of 120 while *Vocanga africana* had the least dominance value of 2, *Elias guinensis* also had the highest importance value indices of 30.3 among the plant species while *Garcinia cola* has the least importance value indices (IVI).

From Table 2 below, shrubs were mostly found in the fallow areas, followed by forested areas and very few were found in the current usage farm site. *Vernonia amygdalina* had the highest abundance value of 94.6% and very high important value indices (IVI) of 130.9, followed by *Manihot esculenta* with abundance value of 31% and important value indices of 17%. *Syzygium samarangens* had the least dominance value of 1% and important value indices (IVI) of 2.3.

The abundance value for forb species showed that *Ipomea aquarica* scored the highest abundance value of 22.2, while *Boreria verticulata* had the least abundance value of 2.5. The evaluation of the importance value indices (IVI) also shows that *Ipomea aquatica* scored the highest importance value indices (IVI) of 19.1, Followed by *Ageratum conyzoides* with the IVI of 15.2 while *Bryophyllum pinnatum* had scored the least (IVI).

Result from the climbers showed that *Telfaria occidentalis* followed by *Peuraria phaseoloides* scored the highest abundance of while *Cissus araloides* and *T.occidentalis*, had the highest importance value indices (IVI) of 35 and 33.5. Also *Cucurbita pepo* score the least abundance value of 3.5 and the least IVI value of 13.1.

Table 3 shows the result for the grass species. *Z. mays* scored the highest abundance and IVI of 58.3 and 52.8 respectively while *Seteria babarta* had the least abundance and (IVI) of 3.5 and 3.9. Also from the graphical analysis showing the IVI of the grass species as presented in Figure 4.5a and b. *Zea mays* was also found to have the highest importance value index.

In the study, the assessment of specie composition of Neni-Nimo degraded watershed, it was discovered that trees were the most dominant in the upper catchment area, and the most frequent tree species is *Elaeis guinensis*, (55%) however, the most dominant tree species also was *Elaeis guinensis*. For the shrubs, *Vernonia amygdalina* was the most abundant and the most dominant specie, with IVI of 56.8. At the middle catchment, shrubs were the most abundant with the most dominant specie being *Chromolaena odorata*, with abundance of 9.5 and Importance value indices IVI of 15.1. Also at the lower catchment area, shrubs were the most abundant, followed by grasses, *Vernonia amagdalaena* had the highest abundance of 49.9 for shrubs, followed by *Zea mays*, with abundance of 28.3 for grasses.

The study revealed that most of the dominant species in the three catchment areas were mostly economic crops which are planted by man.

**Table 1:** Importance value indices and the dominance for trees

Species	<i>Albizia Lebeck</i>	<i>Albizia chaeverieri</i>	<i>Afzelia Africana</i>	<i>African elemi</i>	<i>Annona muricata</i>	<i>Azadiracta indica</i>	<i>Citrus sinensis</i>	<i>Brachystegia eurycoma</i>	<i>Bridelia snigeritiana</i>	<i>Bridelia ferruginea</i>	<i>Casuarina equisetifolia</i>	<i>Dactydenia barteri</i>	<i>Detarium microcarpum</i>	<i>Diallum guineensis</i>	<i>Elaeis guineensis</i>	<i>Erythrophleum suaveolens</i>	<i>Gmelina arborea</i>	<i>Garcinia cola</i>
ARI	2	2		1	1			1			2	2	2	2		1	4	
AR2	1	2	2	1					1	1	1			4	10	4	5	1
AR3			1	2		2				2	1	3	1	3	7		3	
BR1							2								5			
BR2						1	1								6			
BR3						2									8			
CRI																		
CR2																		
CR3																		
Total	3	4	3	4	1	5	3	1	1	3	4	5	3	9	36	5	12	1
Density	0.3	0.4	0.3	0.4	0.1	0.5	0.3	0.1	0.1	0.3	0.4	0.5	0.3	1	4	0.5	1.3	0.1
Frequency	22.2	22.2	22.2	33.3	11.1	33.3	22.2	11.1	11.1	22.2	33.3	22.2	22.2	33.3	55.5	22.2	33.3	11.1
Rel. density	1.5	1.9	1.5	1.9	0.5	2.5	1.5	0.5	0.5	1.2	1.9	2.5	1.5	4.9	19.6	2.5	6.3	0.5
Rel. Frequency	2.173	2.173	2.173	3.26	1.09	3.26	2.173	1.09	1.09	2.173	3.26	2.173	2.173	3.26	5.43	2.173	3.26	1.09
$\bar{x}$ GBH	45	37	22	26	62	21.2	28.7	154.2	55.4	28.0	84.0	48.0	24.3	32.0	32.0	51.4	195.6	46.5
Basal area	11.04	7.47	2.64	3.69	20.9	2.4	4.5	129.6	16.7	4.27	38.5	12.57	3.14	5.6	9.6	14.4	6.5	11.8
Dominance	17.6	15.8	4.2	7.8	11	6.36	7.2	68.7	8.9	6.8	81.62	33	4.9	29.6	120	15.12	27.5	6.3
Rel. dominance	1.8	1.6	0.4	0.8	1.1	0.6	0.7	6.9	0.9	0.7	8.3	3.4	0.5	3.0	12	1.5	2.8	0.6
IVI	5.5	5.7	6.1	5.9	2.69	6.4	4.4	8.5	2.5	4.5	13.6	8.1	4.2	11.2	37	6.2	12.36	2.2

**Table 1:** continuation: showing the importance value indices and the dominance for trees.

Species	<i>Irvingia gabonensis</i>	<i>Mangifera indica</i>	<i>Milicia regia</i>	<i>Milicia excels</i>	<i>Myrianthus arboreus</i>	<i>Nauclear latifolia</i>	<i>Newbouldia leavis</i>	<i>Napolenea imperalis</i>	<i>Pentaclethra macropylla</i>	<i>Psidium guajava</i>	<i>Psorospermum febrifugum</i>	<i>Raphia regalis</i>	<i>Rothmania hispidia</i>	<i>Senna seamea</i>	<i>Sterculia tragacantha</i>	<i>Spondias monbin</i>	<i>Tratinikia rhatifolia</i>	<i>Tetrapluera tetraptera</i>
ARI			1	1	2	1		2			1	4	3	2	1	3		2
AR2					2	2		2	1		2	2	2	1	2	5	3	
AR3							2	3					3			3	2	1
BR1	1	2								4								
BR2		1					4	1										
BR3	2	1					3			2								
CRI																		
CR2																		
CR3																		
Total	3	4	1	1	4	3	9	8	1	6	3	6	8	3	3	11	5	3
Density	0.3	0.4	0.1	0.1	0.4	0.3	1	0.8	0.1	0.6	0.3	0.6	0.8	0.2	0.3	1.2	0.5	0.3
Frequency	22.2	33.3	11.1	11.1	22.2	22.2	33.3	44.4	11.1	22.2	22.2	22.2	33.3	22.2	22.2	33.3	22.2	22.2
Rel. density	1.5	1.9	0.5	0.5	1.9	1.5	4.9	3.9	0.5	2.9	1.5	2.9	3.9	1.5	1.5	5.9	2.5	1.5
Rel. Frequency	2.173	3.26	1.09	1.09	2.173	2.173	3.26	4.26	1.09	2.173	2.173	2.173	3.26	2.173	2.173	3.26	5.43	2.173
$\bar{x}$ GBH	12.0	7.14	109.9	18.3	5.3	3.3	1.3	14.9	172.8	2.6	7.9	3.9	7.1	8.3	2.8	9.2	6.5	31.9
Basal area	47.0	36.2	142	58.0	31.2	24.5	15.2	52.4	206.5	21.7	38.0	27.3	36.4	39.0	22.7	41.2	34.4	76.5
Dominance	19.08	15.13	58.23	9.699	11.24	5.24	6.87	63	91.5	8.24	12.6	12.4	30	13.2	4.3	58	17.2	50.7

Rel. dominance	1.9	1.5	5.9	0.9	1.14	0.5	0.7	6.4	9.2	0.8	1.3	1.3	3.0	1.3	0.5	5.9	1.74	5.2
IVI	10.5	6.7	7.5	2.5	5.2	4.2	9.2	14.6	10.8	5.9	5.9	6.4	10.2	4.9	4.2	15.6	9.7	8.9

**Table 1:** continuation: showing the importance value indices and the dominance for trees continuation

Species	<i>Treculia Africana</i>	<i>Terminalia avicennoides</i>	<i>Voacanga Africana</i>	<i>Zanthaxylon zanthaxyloides</i>	Total
ARI	2	2	1	1	
AR2		3	3		
AR3		2		2	
BR1	1				
BR2	3				
BR3					
CRI					
CR2					
CR3					
Total	6	7	4	3	
Density	0.6	0.7	0.4	0.3	20.4
Frequency	33.3	33.3	22.2	22.2	1021.2
Rel. density	2.9	3.4	1.9	1.5	
Rel. Frequency	3.26	3.16	2.173	2.173	
GBH	24	17.5	14.7	19.2	
Basal area	3.1	1.6	1.2	2.0	
Dominance	9.8	5.9	2.5	3.2	984
Rel. dominance	0.9	0.6	0.2	0.3	
IVI	7.1	7.2	4.3	3.9	

**Table 2:** Showing the important value indices for shrubs

Species	<i>Alchonea cordifolia</i>	<i>Angylocalyx olligophyllus</i>	<i>Annona senegalensis</i>	<i>Calluna vulgaris</i>	<i>Carica papaya</i>	<i>Cassia occidentalis</i>	<i>Chromolaena odorata</i>	<i>Combretum micranthum</i>	<i>Dichrostachys cinerea</i>	<i>Euclinia longifolia</i>	<i>Griffonia simplicifolia</i>	<i>Guinea senegalensis</i>	<i>Manihot esculentum</i>	<i>Mimosa invisa</i>	<i>Occimum sanctum</i>
AR1				19			2			1					
AR2	2		3	24			3		1	1					1
AR3	1		1	16		3	13				7	1			U
BR1	3	2	4	7		15	22	4	2	1		2		2	
BR2		2			1	9	15	7			5			5	2
BR3	2	3	2		3	7	2	12	1		9				3
CRI					2									42	
CR2					3									29	
CR3		1												22	
Total	8	8	10	66	9	34	57	23	4	3	21	3	93	7	6
Density	0.8	0.8	1.1	7.3	1	3.7	6.3	2.5	0.4	0.3	2.3	0.3	10.3	0.7	0.6
Frequency	44.4	44.4	44.4	44.4	44.4	44.4	66.6	33.3	33.3	33.3	33.3	22.2	33.3	22.2	33.3
Rel. density	0.9	0.9	1.2	7.8	1	3.9	6.7	2.7	0.4	0.3	2.4	0.3	10.6	0.7	0.6
Rel. Frequency	3.5	3.5	3.5	3.5	3.5	3.5	5.2	2.6	2.6	2.6	2.6	1.8	2.6	1.8	2.6



Abundance	2	2	2.5	16.5	2.3	8.5	9.5	7.6	1.3	1	7	1.5	31	3.5	2
Rel. abundance	0.7	0.7	0.8	5.4	0.8	2.8	3.2	2.6	0.4	0.3	2.3	0.4	10.4	1.2	0.7
IVI	5.1	5.1	5.5	16.7	5.3	10.2	15.1	7.9	3.4	3.2	7.3	2.5	23.6	3.7	3.9

**Table 2:** continuation: showing important value indices for shrubs.

Species	<i>Ocimum tenuiflorum</i>	<i>Ocimum gratissimum</i>	<i>Olox viridis</i>	<i>Olox subscorpioidea</i>	<i>Phaseolus vulgaris</i>	<i>Piliostigma thonningii</i>	<i>Rauvolfia vomitoria</i>	<i>Sacocephalum laxiflora</i>	<i>Senna alata</i>	<i>Senna senguana</i>	<i>Senna tora</i>	<i>Solanum melanguena</i>	<i>Syzygium samarangens</i>	<i>Uvaria chamae</i>	<i>Vernonia amygdalina</i>	Total
AR1			2			2	1	1		2		2				
AR2				2			1	3			1					
AR3			7		3	2	2	5	1	1	2			2		
BR1	3			5	5		5		3	2	6	11		4	2	
BR2	3		3	4	5	3	3		2	7	9	7	2	7	5	
BR3	2		5	7	7	5	2			5	3	4		9	7	
CRI		3		1		2							1		92	
CR2		5													87	
CR3		4													105	
Total	8	12	17	19	20	14	14	9	6	19	21	24	3	22	298	
Density	0.8	1.3	1.8	2.1	2.2	1.5	1.5	1	0.6	2.1	2.3	2.6	0.3	2.4	33.1	94
Frequency	33.3	33.3	44.4	55.5	44.4	55.5	66.6	33.3	33.3	55.5	55.5	44.4	22.2	44.4	66.6	1265.4
Rel. density	0.9	1.4	1.9	2.2	2.3	1.6	1.6	1	0.7	2.2	2.4	2.7	0.4	2.5	35.2	
Rel. Frequency	2.6	2.6	3.5	4.3	3.5	4.3	5.2	2.6	2.6	4.3	4.3	6	1.8	3.5	5.2	
Abundance	2.6	4	4.3	3.8	5	2.8	2.3	3	2	3.8	4.2	1.7	1	5.5	49.9	297.9
Rel. abundance	0.9	1.3	1.4	1.3	1.7	0.9	0.8	1	0.7	1.3	1.3	1.3	0.3	1.8	16.4	
IVI	4.4	5.3	6.8	7.5	7.5	6.8	7.6	4.6	4	7.8	8	10	2.5	7.8	56.8	

**Table 3:** showing the important value indices of grass species.

Species	<i>Acroceras zizanioides</i>	<i>Andropogon gayanus</i>	<i>Andropogon tectorum</i>	<i>Axonopus compressus</i>	<i>Bambusa vulgaris</i>	<i>Brachiaria deflexa</i>	<i>Brachiaria ruziziensis</i>	<i>Chloris pilosa</i>	<i>Cynopogon citratus</i>	<i>Cynodon dactylon</i>	<i>Hackelochloa granularis</i>	<i>Digitaria horizontalis</i>	<i>Digitaria nuda</i>	<i>Echinochloa obtusiflora</i>
AR1					7						4			7
AR2	5	2	5		5		3			4		4		
AR3	5	5	2		3		4							
BR1	11	11			11	5	4	7			5	20		2
BR2		7				2		2			7	13	7	5
BR3	10	13		7						11	9	7	10	
CRI				4					4	3				
CR2									2	2				
CR3														
Total	31	50	7	11	26	7	11	9	6	20	25	44	17	14

Density	3.4	5.5	0.7	1.2	2.8	0.7	1.2	0.9	0.6	2.2	2.7	4.8	1.8	1.5
Frequency	33.3	55.5	22.2	22.2	44.4	22.2	33.3	22.2	22.2	44.4	44.4	44.4	22.2	33.3
Rel. Density	5	8.2	1	1.8	4.2	1	1.8	1.3	0.9	3.3	4	7.2	2.7	2.2
Rel. Frequency	3.7	6.2	2.5	2.5	4.9	2.5	3.7	2.5	2.5	4.9	4.9	4.9	2.7	3.7
Abundance	10.3	10	3.5	5.5	8	3.5	6.7	4.5	3	6.3	6.3	11	8.5	4.7
Rel. Abundance	4.9	4.8	1.7	2.6	3.8	1.7	3.2	2.2	1.4	3	3	5.2	4	2.3
IVI	13.6	19.2	5.2	6.9	12.9	5.2	8.7	6	8	11.2	11.9	17.3	9.4	8.2

**Table 3:** continuation: showing the important value indices of grass species.

Species	<i>Eleusine indica</i>	<i>Eragrostis alrovirens</i>	<i>Imperata cylindrical</i>	<i>Sorghum arundinaceum</i>	<i>Panicum mmaximum</i>	<i>Panicum repens</i>	<i>Peniserum pepureum</i>	<i>Pennisetum pedicclatitm</i>	<i>Pernnisetum polystachion</i>	<i>Rottboelia cochinchinensis</i>	<i>Setaria barbata</i>	<i>Setaria longiseta</i>	<i>Zea mays</i>	Total
AR1			4				2							
AR2		3	3		9		4			2	5	5		
AR3		4	2		13			7		5		2		
BR1		21	10	11	11			9	7	8				
BR2		11	14	7			11		4	11				
BR3	14		9	9	5	8	7							
CRI					2									25
CR2														28
CR3							3							32
Total	14	39	42	27	40	8	27	16	11	26	5	7	85	
Density	1.5	4.3	4.6	3	4.4	0.8	3	1.7	1.2	2.8	0.5	0.7	9.4	67
Frequency	11.1	44.4	66.6	33.3	55.5	11.1	55.5	22.2	22.2	44.4	11.1	22.2	33.3	898.3
Rel. Density	2.2	6.4	6.9	4.5	6.6	1.2	4.5	2.5	1.8	4.2	0.7	1	14	
Rel. Frequency	1.2	4.9	7.4	3.7	6.2	1.2	6.2	2.5	2.5	4.9	1.2	2.5	3.7	
Abundance	14	9.8	7	9	8	8	5.4	8	5.5	6.5	5	3.5	28.3	208.8
Rel. Abundance	6.7	4.7	3.4	4.3	3.8	3.8	2.5	3.8	2.6	3.1	2.3	1.7	13.6	
IVI	10.1	16	17.7	12.3	16.6	6.6	13.2	8.8	6.9	12.2	4.2	4.5	31.3	



## Conclusion

In conclusion, the studies on specie composition and abundance of Neni-NImo watershed, shows that that, anthropogenic influence, has led to the drastic reduction in abundance of plants composition of the watershed, and the most dominant species are those economic crop cultivated by man.

## References

1. Arrhenius O. Species and Area. British Ecological Society, 1921:9(1):95-99.
2. Bhadra AK, Pattanayak SK. Dominance is more justified than abundance to calculate important value indices (IVI). Asian Journal of Science and Technology, India, 2017:7(9):3577-3601.
3. Boyer EW, Goodale CL, Jaworsk NA, Howarth RW. Anthropogenic nitrogen sources and relationships to riverine nitrogen export in the northeastern USA. Biogeochemistry, 2002:57(1):137-169.
4. Cottam, G. and Curtis, J.T. Trees: A method for making rapid survey of woodland by means of pairs of randomly selected. Washington D.C. Ecology, 1994:30:101-104.
5. Cox, G.W. Laboratory Manual of General Ecology. 2<sup>nd</sup> Edition. W.M.C. Brown Company Publishers. U.S.A, 1976, 657.
6. Enwelu A, Agwu AE, Igbokwe EM. Challenges of participatory approach to watershed management in rural communities of Enugu State. Nigeria. Journal of Agriculture Extension, 2014:14(1):69-79.
7. Foley JA. Global consequences of land use, Science, California, 2005:309(734):570-574.
8. Hale RL, Hoover JH., Wollheim WM, Vörösmarty CJ. History of nutrient inputs to the northeastern United States, 1930–2000. Global Biogeochemistry Cycles, U.S.A, 2013:27:578-591.
9. Howarth R, Swaney D, Billen G, Garnier J, Hong C, Humborg B *et al.* Nitrogen fluxes from the landscape are controlled by net anthropogenic nitrogen inputs and by climate. Frontier Ecology Environment, U.S.A., 2012:10(1):37-43.
10. Kunstler G, Curt T, Lepart J. Spatial pattern of beech (*Fagus sylvatica* L.) and oak (*Quercus pubescens* Mill) seedlings in natural pine (*Pinus sylvestris* L.) woodlands. Eur Journal Forest Research, 2004:123:331-337.
11. Mishra, R.. Ecology workbook. Oxford and IBH publishing company, Calcuta India, 1968, 242.
12. Okereke CN, Mbaekwe E, Nnabude P, Ekwealor KU, Godwin NN, Chisom FI *et al.* Comparative evaluation of specie richness and diversity of three parallel ecosystem, in South East Nigeria. JAERI, 2016: 8(4):1-12.
13. Philips, E.A. Method of Vegetation Studies. Henry Holt publishers, New York, U.S.A, 1959, 14.
14. Russell MJDE, Weller TE, Jordan KJ, Sigwart KJ, Sullivan KJ. Net anthropogenic phosphorus inputs: Spatial and temporal variability in the Chesapeake Bay Region, U.S.A. Biogeochemistry, 2008:88(3):285-304.
15. Vitousek PM, Mooney HA, Lubchenco J, Melillo J. Human domination of Earth's Ecosystems. Science, 1997:277(5325):494-499.