



International Journal of Biology Sciences

ISSN Print: 2664-9926
 ISSN Online: 2664-9934
 Impact Factor: RJIF 5.45
 IJBS 2022; 4(2): 186-189
www.biologyjournal.net
 Received: 22-09-2022
 Accepted: 24-10-2022

Pragada Vyas
 M.Sc. Scholar, Department of
 Entomology, SHUATS,
 Prayagraj, Uttar Pradesh,
 India

Anoorag R Tayde
 Assistant Professor,
 Department of Entomology,
 SHUATS, Prayagraj, Uttar
 Pradesh, India

Corresponding Author:
Pragada Vyas
 M.Sc. Scholar, Department of
 Entomology, SHUATS,
 Prayagraj, Uttar Pradesh,
 India

Comparative potency of chemical insecticides and biopesticides in management of shoot and fruit borer [*Leucinodes orbonalis* (Guenee)] in Brinjal [*Solanum melongena* (L.)]

Pragada Vyas and Anoorag R Tayde

DOI: <https://doi.org/10.33545/26649926.2022.v4.i2c.121>

Abstract

The present experiment was carried out to compare the efficacy against *Leucinodes orbonalis* and its effect on yield of Brinjal. The best and most economical treatment was Chlorantraniliprole @ 18.5 EC (1:10.2) which was on par with Emamectin benzoate @ 5 SG (1:9.8) followed by Carbosulfan 25 EC (1:7.3), *Beauveria bassiana* (1:6.4) Spinosad @45% SC (1:5.5), followed by *Bacillus thuringiensis* (1:5.2) and Neem oil (1:4.3) as compared to control (1:3.1). The data on the percent infestation of the shoot and fruit borer on brinjal 3rd, 7th, and 14th day after the first spray reveal that all the chemical treatments were significantly superior over control. Chlorantraniliprole @ 18.5 EC (5.94) was par with Emamectin benzoate @5 SG (8.49) followed by Carbosulfan 25 EC (10.71), *Beauveria bassiana* (12.91) Spinosad @45% SC (13.13), followed by *Bacillus thuringiensis* (14.85) and, Neem oil (16.87). The yield among the treatment were significantly higher in Chlorantraniliprole @ 18.5 EC (207.50q/ha) followed by Emamectin benzoate @ 5 SG (183.83q/ha)

Keywords: Biorationals, chlorantraniliprole, cost benefit ratio, chemical insecticide, *Leucinodes orbonalis*

Introduction

Brinjal (*Solanum melongena* Linnaeus) also known as eggplant is referred as “King of vegetables”. It is also called as brinjal in India, and Auvergne in Europe. It belonging to the family “Solanaceae”, the name eggplant derives from the shape of the fruit of some varieties, which are white and shape very similarly to chicken eggs.

Among solanaceous vegetables, brinjal (*Solanum melongena* L.) is one of the important crops grown throughout India. Eggplant is adapted to a wide range of climatic conditions, such as high rainfall and high temperatures from North to South and West to East. It is also one among the few vegetables capable of high yields in hot-wet environments. Brinjal fruits are reasonable sources of vitamins and minerals. (Tripura *et al.*, 2017) [19].

It is the most important vegetable in the Indian Subcontinent that accounts for almost 50% of the world's area under its cultivation. Under sustainable farming, brinjal provides regular daily income to meet the day-to-day expenditure. (Murugesan, 2009) [10] It has become an important economic source for farmers and field laborers.

It is also used as a raw material in pickle making and as an excellent remedy for curing diabetes. It is also used as a good appetizer. It is a good aphrodisiac, cardiotoxic, laxative and reliever of inflammation (Shridhara., 2019) [18].

The estimated total world production for eggplants in 2020 was 56,618,843 metric tons, up by 2.2% from 55,376,521 tons in 2019 China is leading in production with 32.03 million tons, the area with and 0.78 million ha, and productivity with 40.96 tons per ha, respectively in the whole world during the year 2016- 17. India is the second largest producer of brinjal being cultivated over an area of 749,000 (ha), production of 12874,000 (MT) with an average annual production of 17.5 million tons per ha in the year 2017-18. In India, it is widely grown in West Bengal, Odisha, Gujarat, Madhya Pradesh, Chhattisgarh, Maharashtra, Andhra Pradesh, Haryana, Assam, Uttar Pradesh, Jharkhand, and Tamil Nadu. Brinjal crop is cultivated in Uttar Pradesh over an area of 8.82, 000ha with an annual production of 275.40 (thousand tons) and productivity of 34.40 (MT/ha) in the year 2020-21, while West Bengal

ranks 1st having the area and production of 163.15 (ha) and 3027.00 (million tons), respectively collection.

in the year 2017-18. However, U.P. is at the apex in productivity over India. It harbors more than 140 species of insect pests belonging to 50 families from 10 orders reported on the crop of the world. (Chowdury *et al.*, 2017).

There are 26 insect-pest species and few no insect-pests species infesting brinjal of which the shoot and fruit borer, (*Leucinodes orbonalis* Guen.); Budworms, (*Scrobipalpa blasigona*;) whitefly, (*Bemisia tabaci* Genn.) leafhopper, (*Amarasca devastans*) (Distant); jassid, *A. (biguttula biguttula* Ishida) epilachna beetle, (*Henosepilachna vigintioctopunctata*) Fab.; aphid, (*Aphis gossypii* Glover.); mealy bug, (*Centroccocus insolitus* Guen), lace wing bug, (*Urentiushy stricellus* Richt). and non-insect pest, red spider mite, (*Tetranychus macfurlanei*) Andre cause severe damage, necessitating initiation of control measures quite frequently (Srinivasan, 2009) of these, the brinjal shoot and fruit borer is considered the main constraint as it damages the crop throughout the year. This pest reported from all brinjal growing area of the world. It is known to damage shoot and fruits of brinjal in all stages of its growth. The yield loss by this pest varied from 0.081.11 q/ha based on the inconsumable pest of damaged fruits and 0.46- 3.80 q/ha when the whole of damaged fruits was taken into consideration.

The Larvae of this pest cause 12-16% damage to shoots and 20-60% damage to fruits. The pest is very active during the rainy and summer season and often causes more than up to 95% in India. It is also reported that the infestation of fruit borer causes a reduction in Vitamin "C" content to an extent of 68 % in the infested fruits. Sarkar *et al.*, (2002) The yield loss in brinjal due to the pest complex is to the extent of 70-80 per cent. It is estimated that the economic injury level equals to 6% infestation of shoot and fruit in India. (Anwar *et al.*, 2015)^[2]

Materials and Methods

The experiment was conducted at the Central Research Farm, Department of Entomology, Nani Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj, Uttar Pradesh, during the *Rabi* season of 2021-22. Prayagraj is situated at an elevation of 78 meters above sea level at 25.87 North latitude and 81.15° E longitudes.

Preparation of Insecticidal spray solution

The desired concentration of insecticidal spray solution of desired concentration for each treatment was freshly prepared each and every time at the site of experiment, just before the start of spraying operations. The quantity of spray materials required for crop was gradually increased as the crop advanced in age.

The spray solution of desired concentration was prepared by adopting the following formula:

$$V = \frac{(C \times A)}{\% a.i.}$$

Where, V=Volume of a formulated pesticide required= Concentration required. A= Volume of total solution to be prepared % i.e., = Given Percentage strength of a formulated pesticide The population of brinjal shoot and fruit borer was recorded before 1- day spraying and on 3rd day, 7th day and

14th day after insecticidal application. The populations of brinjal shoot and fruit borer was recorded on 5 randomly selected and tagged plants from each plot and then it was converted into per cent of infestation by following formula.

On Shoot

At each picking the total number of shoots and number of shoots infested of five selected plants from each treatment replication wise were recorded.

$$\% \text{ Shoot infestation} = \frac{\text{No. of shoot infested}}{\text{Total no. of shoot}} \times 100$$

On Fruit:

At each picking the total number of fruits and number of fruits infested of five selected plants from each treatment replication wise were recorded.

$$\% \text{ Fruit infestation} = \frac{\text{No. of fruit infested}}{\text{Total no. of fruit}} \times 100$$

Results and Discussion

The data on the percent infestation of shoot and fruit borer on brinjal 3rd, 7th and 14th day after first spray revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest per cent shoot, infestation was recorded in Chlorantraniliprole 18.5 SC (6.84), Abamectin benzoate 5 SG (9.78), Carbosulfan 25 EC (12.40), Spinosad 45% SC (13.90), *Beauveria bassiana* (14.68), *Bacillus thuringiensis* (16.23), and Neem oil (19.08). The treatments Neem oil (19.08) was least effective among all the treatments. Control plot (24.31) infestation.

The data on the percent infestation of shoot and fruit borer on brinjal 3rd, 7th and 14th day after second spray revealed that all the chemical treatments were significantly superior over control. Among all the treatments lowest per cent shoot, infestation was recorded in Chlorantraniliprole 18.5 SC (5.05), Emamectin benzoate 5 SG (7.29), Carbosulfan 25 EC (9.03), *Beauveria bassiana* (11.14), Spinosad 45% SC (12.36), *Bacillus thuringiensis* (13.47) and Neem oil (14.67), was least effective among all the treatments. Control plot (28.33) infestation. The treatments Neem oil (14.67) was least effective among all the treatments. Control plot (28.33) infestation.

These results are in support with Tripura *et al.*, (2017)^[19], Reddy and Kumar (2022)^[12], Jat *et al.*, (2020)^[6] reported that the treatment Chlorantraniliprole was superior in reducing the population of shoot and fruit borer. Vinayaka *et al.*, (2019)^[20], Warghat *et al.* (2020)^[21], Goud *et al.*, (2019)^[5] reported that Emamectin benzoate maximum protection and minimum shoot damage in brinjal), Chandar *et al.*, (2020)^[3], Das *et al.*, (2016)^[4], Mahla *et al.*, (2017)^[9], Anand *et al.*, (2014)^[11] also reported that Carbosulfan 25 EC as most effective chemical to control the brinjal shoot and fruit borer. Kumar and Thakur (2017)^[8], Soulakhe *et al.*, (2021)^[17] also reported that next effective treatment in controlling the pest population of shoot and fruit borer. Sureshsing and Tayde (2017)^[16], Kapuriya *et al.*, (2019)^[7] Jat *et al.*, (2020)^[6] also reported that Spinosad 45% SC was effective in reducing the infestation of shoot and fruit borer, Rashid *et al.*, (2018)^[11], Singh *et al.*, (2016)^[15] reported that *Bacillus thuringiensis* SC was also effective in reducing the infestation of shoot and fruit borer Sangma *et al.*, (2019)^[13], Murugesan and Murugesh (2009)^[10], also reported that

Neem oil is effective in reducing the infestation of shoot and fruit borer.

The yields among the treatment were significant. The highest yield was recorded in Chlorantraniliprole 18.5 SC (207.50 q/ha) these findings were supported by Jat *et al.*, (2020) [6], Emamectin benzoate 5 SG (183.83 q/ha) these findings were supported by Warghat *et al.* (2020) [21], Carbosulfan 25 EC (137.9 q/ha) these findings were supported by Mahla *et al.*, (2017) [9], *Beauveria bassiana* (124.6 q/ha) Soulakhe *et al.*, (2021) [17], Spinosad 45% SC (106.5 q/ha), *Bacillus thuringiensis*

(98.3 q/ha) Jat *et al.*, (2020) [6] and Neem oil (87 q/ha) Sangma *et al.*, (2019) [13]. The treatments Neem oil (87 q/ha) was least effective among all the treatments. Control plot (56.6 q/ha) yield. Cost benefit ratio revealed, interesting result such as, Chlorantraniliprole 18.5 SC (1:10.8) the best and most economical treatment followed by Emamectin benzoate 5 SG (1:9.8), Carbosulfan 25 EC (1:7.3), *Beauveria bassiana* (1:6.4), Spinosad 45% SC (1:5.5), *Bacillus thuringiensis* (1:5.2), and Neem oil (1:4.3) and control (1:3.1).

Table 1: Efficacy of biopesticides and chemicals on damage of shoot and fruitborer, *Leucinodes orbonalis* on brinjal.

	Per cent shoots and fruit infestation of <i>Leucinodes orbonalis</i>	First spray								Second spray			Yield	B:C
		1DBS	3 DAS	7 DAS	14DAS	Mean	3DAS	7 DAS	14DAS	Mean				
T ₁	<i>Beauveria bassiana</i> 1% WP (1x10 ⁸ CFU/ml)	20.80	14.91	11.44	17.69	14.68	11.63	09.17	12.63	11.14	124.6	1:6.6		
T ₂	<i>Bacillus thuringiensis</i> (1x10 ⁹ CFU/ml)	20.24	19.10	13.50	16.10	16.23	13.50	12.35	14.57	13.47	98.30	1:5.1		
T ₃	Emamectin benzoate @ 5% SG	20.81	12.51	07.33	09.52	09.79	07.33	05.18	09.38	07.20	183.83	1:9.8		
T ₄	Spinosad 45 SC	20.97	16.02	12.49	13.21	13.90	12.39	11.15	13.54	12.36	106.5	1:5.5		
T ₅	Carbosulfan 25 EC	19.77	13.26	09.38	14.56	12.40	08.37	08.31	10.39	09.02	137.9	1:7.3		
T ₆	Chlorantraniliprole 18.5 SC	20.24	11.20	03.37	05.96	06.84	0.19	02.22	06.75	05.05	207.50	1:10.8		
T ₇	Neem oil @2%	20.70	20.71	14.69	21.85	19.08	14.69	13.50	15.84	14.67	87	1:4.3		
T ₈	Control	20.41	24.22	22.98	25.73	24.31	28.30	27.07	29.63	28.33	56.6	1:3.1		
	Overall Mean	20.49	16.48	11.89	15.57	14.65	12.08	11.11	14.09	12.65	-	-		
	F- test	NS	S		S	S	S	S	S	S	-	-		
	S. Ed. (±)	00.87	01.02	01.13	01.29	01.39	01.50	01.40	01.20	00.52	-	-		
	C. D. (P = 0.05)	01.86	02.22	04.25	02.77	02.99	03.24	03.04	02.63	01.14	-	-		

Conclusion

From the present study, the results showed that chlorantraniliprole @18.5 SC is most effective treatment against brinjal fruit and shoot borer producing maximum yield and recorded highest Cost-Benefit ratio compared to other treatments. While emamectin benzoate 5% SG and carbosulfan 25 EC has shown average results. Spinosad 45 %SC and *Beauveria bassiana* and *Bacillus turingiensis* has proved to be least effective chemicals. Botanical neem oil 2% found to be least effective in managing *Leucinodes orbonalis*. Botanicals are the part of integrated pest management in order to avoid indiscriminate use of pesticides causing pollution in the environment and not much harmful to beneficial insects.

References

- Anand GS, Sharma RK, Shankarganesh K. Evaluation of bio-efficacy and compatibility of emamectin benzoate with neem based biopesticide against fruit borers of brinjal and okra. Indian Journal of Agricultural Sciences. 2014;84(6):746-53.
- Anwar S, Mari JM, Khanzada MA, Ullah F. Efficacy of insecticides against infestation of brinjal fruit borer, (*Leucinodes orbonalis* (Guenee) (Pyralidae: Lepidoptera) under field conditions. Journal of Entomology and Zoology Studies. 2015;3(3):292-295.
- Chandar AS, Kumar A, Singh U, Kakde AA, Nawale JS, Narode MK *et al.* Efficacy of certain chemicals and biopesticides ag ainst brinjal shoot and fruit borer *Leucinodes orbonalis* (Guenee). Journal of Entomology and Zoology Studies. 2020;8(5):220-223.
- Das G. Comparative bioefficacy of different insecticides against fruit and shoot borer (*Leucinodes orbonalis* (Guenee) of brinjal and their effect on natural enemies. International Journal of Green Pharmacy (IJGP). 2016;10(04).
- Goud GS, Bondre CM, Gawali KA. Efficacy of some chemical and botanical pesticides against brinjal shoot and fruit borer *Leucinodes orbonalis* (Guenee). Journal of Pharmacognosy and Phytochemistry, 2019;8(4):2453-2455.
- Jat HK, Shrivastava VK, Dubey R. Management of the brinjal shoot and fruit borer *Leucinodes orbonalis* (Guen.) through newer insecticides in brinjal. Journal of Entomology and Zoology Studies. 2020;8(5):1259-1261.
- Kapuriya H. Periodical occurrence and effectiveness of selected insecticides in regulating brinjal shoot and fruit borer (*Leucinodes orbonalis* (Guenee). Journal of Pharmacognosy and Phytochemistry. 2019;8(1):2314-2316.
- Kumar A, Thakur S. Comparative efficacy of essential oils, neem products and *Beauveria bassiana* against brinjal shoot and fruit borer (*Leucinodes orbonalis* (Guenee) of Brinjal (*Solanum melongena* L.). Journal of Entomology and Zoology Studies. 2017;5(4):306-309.
- Mahla MK, Kumar A, Lekha AV, Singh V. Bio-efficacy of carbosulfan 25 EC against fruit and shoot borer of brinjal and their natural enemies. Journal of Entomology and Zoology Studies. 2017;5(5):260-264.
- Murugesan N, Murugesan T. Bioefficacy of some plant products against brinjal fruit borer, *Leucinodes orbonalis* (Guenee) (*Lepidoptera: Pyralidae*). Journal of Biopesticides. 2009;2(1):60-63.
- Rashid MA, Hasan MK, Matin MA. Socio-economic performance of Bt eggplant cultivation in Bangladesh. Bangladesh Journal of Agricultural Research. 2018;43(2):187-203.
- Reddy CSTS, Kumar A. Efficacy of selected insecticides against brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee). The Pharma Innovation Journal. 2022;11(4):1327-1330.

13. Sangma CD, Simon S, Nagar S. Indigenous pest control practices for the management of Brinjal shoot and fruit borer (*Leucinodes orbonalis* (Guen). Journal of Pharmacognosy and Phytochemistry. 2019;8(3):4221-4223.
14. Paswan A, Choudhary AS, Raj S, Sonloi P, Sonwani A. Effect of integrated nutrient management on yield of Brinjal. Int. J Agric. Food Sci. 2022;4(1):12-16. DOI: 10.33545/2664844X.2022.v4.i1a.59
15. Singh JP, Gupta PK, Chandra U, Singh VK. Bioefficacy of newer insecticides and biopesticides against brinjal shoot and fruit borer *Leucinodes orbonalis* (Guenee) (Lepidoptera: Pyralidae). International Journal of Plant Protection. 2016;9(1):1-7.
16. Sureshsing MC, Tayde AR. Efficacy of certain biorationals against shoot and fruit borer (*Leucinodes orbonalis* (Guenee) of brinjal (*Solanum melongena* L.). Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1857-1859.
17. Soulakhe AB, Bantewad SD, Jayewar NE. Biorational management of brinjal shoot and fruit borer *Leucinodes orbonalis*. The Pharma Innovation Journal. 2021;10(8):617-623.
18. Shridhara M, Hanchinal SG, Sreenivas AG, Hosamani AC, Nidagundi JM. Evaluation of newer insecticides for the management of brinjal shoot and fruit borer *Leucinodes orbonalis* (Guenee) (Lepidoptera: Crambidae). International Journal of Current Microbiology and Applied Sciences. 2019;8(3):2582-2592.
19. Tripura A, Chatterjee ML, Pande R, Patra S. Biorational management of brinjal shoot and fruit borer (*Leucinodes orbonalis* (Guenee)) in mid hills of meghalaya. Journal of Entomology and Zoology Studies. 2017;5(4):41-45.
20. Vinayaka KS, Singh B, Yadav SS, Nayak SB. Efficacy of Different Insecticides against brinjal fruit borer (*Leucinodes orbonalis* (Guenee) and their impact on fruit yield. Journal of Entomology and Zoology Studies. 2019;7(3):66-66.
21. Warghat AN, Nimbalkar D, Tayde AR. Bio-efficiency of some insecticides against Brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guen). Journal of Entomology and Zoology Studies. 2020;8(1):932-936.
22. Sarker M, Uddin MM, Howlader MTH. Management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee) using biorational insecticide based IPM packages. Fundamental and Applied Agriculture. 2020;5(3):361-371.