

ISSN Online: 2664-9934 Impact Factor: RJIF 5.45 IJBS 2023; 5(2): 19-21 www.biologyjournal.net Received: 15-05-2023 Accepted: 21-06-2023

ISSN Print: 2664-9926

Sanjay Prasad Gupta National Research Laboratory for Concernation of Cultural

for Conservation of Cultural Property, Lucknow, India

Sachin Kumar Agnihotri

Archaeological Survey of India, Science Branch, Agra, India

Fungal deterioration of historical monument with reference to Jaleshwarnath temple of Shivpur, Chhattisgarh, India

Sanjay Prasad Gupta and Sachin Kumar Agnihotri

DOI: https://dx.doi.org/10.33545/26649926.2023.v5.i2a.166

Abstract

Stone objects may support novel communities of microorganisms that are active in bio-deterioration process. Bio-film on the sandstone monuments contains a complex of consortia of Bryophyta and Fungi. The Bryophyta make up the photosynthetic part of the bio-film while hyphae, filaments and spores take part as fungal components. These structures make a dense layer by intertwining over the surface of sandstone monuments. In the present investigation 10 samples were collected from different sites of Jaleshwarnath temple of Shivpur of Chhattisgarh and their surroundings. The five (05) fungal species were isolated which have dominance over sandstone structures of the monuments. During investigation period it was observed that *Aspergillus sp.* was found most dominant. The identified micro fungi cause discoloration as well as mechanical exfoliation of building stone material that was analyzed through mechanical hyphae penetration and production of dark pigments and organic acids.

Keywords: Degradation, deterioration, sandstone monument, micro-flora, bio-film, hyphae, discoloration, dominance etc.

Introduction

Shivpur is a little town situated around 40 kilo meter far from district head quarter of Surguja of Chhattisgarh state in India. This antiquated sanctuary devoted to Lord Shiva is situated in Shivpur town in the midst of pleasant characteristic environment (under the sky). This sanctuary is otherwise called Jaleshwarnath Mandir. It is a mainstream journey goal that draws in an immense horde of devotees from distant spots. On the event of Maha Shivratri and Basant Panchami, an enormous celebration is arranged here. This ancient temple is built in Shivpur village near Pratappur of Chhattisgarh. This temple is the most famous and beautiful place of Pratappur of Surguja district. There is dense forest and mountains all around here. This temple is located a little away from Pratappur-Balrampur Highway Road. Here Shivling is sitting in the main sanctum sanctorum. The specialty of this temple is that Shivling is sitting here. Here water flows from the sitting point of Shivling and this water continues to flow like this throughout the year and for 24 hours. There is a water kund is made here, where lotus flowers are planted. This temple is very well built. A fair is organized here on Sawan Monday and Mahashivaratri, in which people from far and wide come to visit. Many deities are available here for darshan. Here Nandi Bhagwan, Jagannath Puri, huge idol of Hanuman Ji, ancient Shivling and many more idols are available to darshan. All these images are very beautiful. This place is very beautiful and people will love coming

Corresponding Author: Sanjay Prasad Gupta National Research Laboratory for Conservation of Cultural Property, Lucknow, India




Fig 1: a) Shiv linga situated in main sanctum sanctum, b) Front view of Jaleshwarnath temple, c) Lateral view of temple d) River near the Jaleshwarnath temple




Fig 2: Fungal deteriorated samples, surrounding of temple

Conservation issue

Jaleshwarnath temple built of sand stone which is porous in nature. The stone surface of the temple has become blackish in appearance due to deposition of dist, dirt, and dried vegetation and micro vegetation growth on an exterior as well as interior portions. The depositions are seeming to be very old as depicted by the formation of secondary lichens of dull green pale white colour which are present all over the stone surface. Due to these depositions the aesthetic beauty of the temples is affected badly. Furthermore, scientific point of view these depositions are very harmful for the health of the stone surface because these microorganism secrets acid which dissolves the component of sand stone. Micro-organism growth such as algae, fungus and lichen on stone surfaces had affected the structure badly, developing cracks as well as porosity of stones.

Increase in humidity level due to high rainfall and also thick forest cover had affected the stone surface. The exterior portion of temple had become blackish due to deposition of dried moss, lichens & bacterial slime. Accumulation of dust and dirt was also a base for the growth of micro vegetation which secretes organic acid causing which harmful to the structure (Kumar & Kumar, 1999) [7]. Stone chipping of temple was taking place which was in need of urgent consolidation.

Material and Method

a) Sampling and Isolation of Fungi

A total of 10 Samples were collected from various locations of Jaleshwarnath temple & surrounding of the temple and brought to the laboratory under aseptic conditions. The isolation of micro- organisms was done by culturing the

samples and by direct incubation of samples in moist chamber. The purified fungal cultures were identified by using mycological techniques and were compared with the available authentic literature, reviews and mycological manuals (Alexopoulus, 1978; Barnett & Hunter, 1978; Gilman, 1995 and Gupta, *et al.*, 2015) [2, 3, 5, 6].

b) Calculations

Various myco-ecological parameters have been calculated using the following formulae:

Frequency (F %) = \frac{Number of plates in which specific organism occurred}{Total number of plates examined} \text{ X 100}

Density (D) = $\frac{\text{Total number of colonies of specific organism}}{\text{Total number of plates examined}}$

 $Abundance (Ab) = \frac{Total\ number\ of\ colonies\ of\ specific\ organism}{Number\ of\ plates\ in\ which\ specific\ organism\ occurred}$

Observation table 1

Isolated fungi	Culture plates										F %	D	A 1.
	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	F %0	ע	Ab
Aspergillus fumigates (Fresenius.)	1	-	-	2	-	-	1	-	-	-	30	0.40	1.33
Aspergillus niger (Tieghem.)	-	6	-	4	1	6	-	-	5	2	60	2.40	4.00
Fusarium oxysporium (Schlechtendahl.)	-	-	8	2	-	-	1	-	-	1	40	1.20	3.00
Rhizopus nodosus (Namyslowski.)	-	-	-	-	2	-	-	-	-	-	10	0.20	2.00
Chaetomium globossum (Kunze.)	-	2	-	-	-	-	-	4	-	-	20	0.60	3.00

Results and Discussion

During screening for search of mycoflora, total five species of fungal organisms were isolated from Jaleshwarnath temple at Shivpur. In the temple Aspergillus niger shows maximum frequency, Density as well as Abundance followed by Fusarium oxysporum& Chaetomium globossum. Some of the fungal species are confined to particular area i.e. Aspergillus fumigates, Chaetomium globossum and Rhizopus nodosus. These confinements of fungal species depend on environmental conditions of the area, which varies from geographical area to area (Gupta et al. 2015) [6]. In the present study Aspergillus species are the most common species found in the sites (Observation table-1).

It has also been shown in the laboratory that fungal species such as *Aspergillus niger* were able to solubilize powdered stone and chelate various minerals in a rich glucose medium because they produce organic acids such as gluconic, citric, and oxalic acids (Lapidi & Schipa, 1973) ^[1]. The toxic metabolites produced by various species of fungal organisms function as chelating agents that can leach metallic cations, such as Iron, Magnesium etc. from the stone surface. Laboratory experiments have demonstrated that basic rocks are more susceptible to fungal attack than acidic rocks. In the present study *Aspergillus* are the most common species found in the sites. *Aspergillus niger* released certain metal ions from the rock samples (Boyle & Voight, 1973 and Gupta, *et al.*, 2015) ^[4,6].

Acknowledgement

Author are gratitude to thanks Dr. Alok Tripathi, Director General, NRLC, Lucknow and Additional Director General, Archaeological Survey of India, New Delhi for their encouragement and guidance.

References

- 1. Lapidi A, Schipa G. Some aspects of the growth of chemotrophic and heterotrophic bacteria from decayed stone" In: Proceedings of the 5th. International Congress on deterioration and conservation of stone. (Ed) G. Felix Lausanre, Switzerland; c1973. p. 633-640.
- Alexopoulos CJ. Introductory mycology (2nd ed.). Wiley Estern Ltd. New Delhi, Banglore, Bombay; c1978.
- Barnett HC, Barry Hunter B. Illustrated genera of important fungi. Macmillan Publishing Company. New

- York and Collier. Macmillon Publishers, London; c1978.
- 4. Boyle JR, Voight GK. Biological weathering of silicate minerals: Implications for tree nutrition and soil genesis. Plant and Soil. 1973;38:191-201.
- 5. Gilman C Joseph. A Manual of Soil Fungi. Print well publication, Jaipur (India); c1995.
- Gupta SP, Sharma DN, Chandrol GK, Misra R, Minj DK. Myco-Ecological Study of Fungal Flora on Deteriorating Sandstone Structures of Historical Monument Kala Dera-I Temple of Jammu & Kashmir State (India), IJIRSET. 2015 Oct;4(10):10298-10302.
- 7. Kumar R, Kumar AV. Biodeterioration of stone in tropical environments USA J Paul Getty Trust. 1999;(0-89236-550):1-2.