



# International Journal of Biology Sciences

ISSN Print: 2664-9926  
 ISSN Online: 2664-9934  
 Impact Factor: RJIF 5.45  
 IJBS 2024; 6(1): 176-182  
[www.biologyjournal.net](http://www.biologyjournal.net)  
 Received: 18-11-2023  
 Accepted: 25-12-2023

**Amanuel Mhreteab**  
 Entomologist, Ministry of  
 Health, Gash Barka Region,  
 Eritrea

**Mensur Yenus**  
 Entomologist, Ministry of  
 Health, Anseba Region,  
 Eritrea

**Awet Mebrahtu**  
 Public Health Officer, Ministry  
 of Health, Gash Barka Region,  
 Eritrea

**Corresponding Author:**  
**Amanuel Mhreteab**  
 Entomologist, Ministry of  
 Health, Gash Barka Region,  
 Eritrea

## First detection of *anopheles stephensi* (Diptera: culicidae) in Gash Barka Region: Eritrea

**Amanuel Mhreteab, Mensur Yenus and Awet Mebrahtu**

**DOI:** <https://dx.doi.org/10.33545/26649926.2024.v6.i1c.203>

### Abstract

**Background:** *Anopheles (An) gambiae* and *(An) funestus* are the dominant vectors in Africa. These vectors inhabit the rural area and as a result malaria was being restricted to the rural areas. Recently, *(An) stephensi* as a competent vector of both *Plasmodium (p). vivax* and *P. falciparum* appear to be better adapted in urban and peri-urban areas of East African countries. So far, *An. Stephensi* found to inhabit in Djibouti, Ethiopia, Sudan, and Somalia and further distributed in Nigeria and Kenya, since 2012. If the invasion of *anopheles stephensi* continue across the African continent, additional 126 million people from urban areas will be at risk of malaria infection. Gash Barka (GB) Region has shared a border with Ethiopia and Sudan (*An. Stephensi* prevalent areas). So the current surveillance aimed to ascertain the presence of *An. Stephensi* in GB region using morphological and molecular identification.

**Method:** Cross sectional entomological survey was conducted during the period Sep. - Dec. 2022 in three towns of Gash Barka Region (*Vis.* Tesseney, Agurdet and Gulj) to assess the presence of *An. Stephensi*.

**Result:** This is the first report in Eritrea to identify the invasive malaria vector *An. Stephensi*. Cement basin was the principal breeding habitat in the study site. It was accountable for more than 90% of the immature anopheles species. Other habitats include plastic/metal basin barrels, clay pots and overhead tanks were found with immature mosquitos.

**Conclusion:** *An. Stephensi* co-exists with native vector species in the study sites. Further research is needed to determine its relative role in malaria transmission in the Region. Health authorities need to revise the existing vector control strategies to target *An. Stephensi* alongside the native malaria vector species.

**Keywords:** WHO, *P. falciparum*, *An. Stephensi*

### Introduction

According to world health organization (WHO) annual report 2021, an impact limitation was occurred in 2015-2020. The progress of malaria control and elimination was remained the same. The report revealed African region as a whole missed the milestone of the Global Technical Strategy for malaria (GTS) 2020, incidence and mortality reduction by 38% and 40%, respectively. In Botswana, Comoros, Eritrea and Madagascar, the malaria mortality rate were increased by 40% or more (WHO, 2021) [32]. In 2021 alone there were 247 million cases and 619,000 deaths globally. The sub Saharan Africa (SSA) were responsible for the 96% of the cases and 96% of deaths (WHO, 2022) [34]. *Anopheles arabiensis*, a member of the *Anopheles gambiae* species complex, is the principal malaria vector in East African countries like Djibouti, Sudan, Somalia, and Ethiopia, as well as in some countries of the Arabian Peninsula, such as Yemen and Saudi Arabia (WHO, 2022) [34]. Twenty years ago, a study conducted in Eritrea investigated the distribution of *Anopheles* mosquitoes throughout the country. The study identified thirteen species, and *Anopheles arabiensis* emerged as the most dominant vector, accounting for more than 80% of the total (Shililu *et al.* 2007) [20]. A similar study conducted in the Anseba region of Eritrea in 2020 also revealed the same result; *Anopheles gambiae* was found to be the dominant vector in that particular location (Yenus *et al.*, 2021) [36]. It is believed that *An. arabiensis* is a rural malaria vector breeds in a variety of breeding habitats like streams, animal hoof prints, river edges, in both temporary and permanent ground pools (Yenus *et al.*, 2021, Shililu *et al.*, 2007, Sinka *et al.*, 2010, Himeidan and Rayah 2008) [36, 20, 24, 16].

In Africa, malaria has been restricted to the rural areas due to the vectors breeding behavior (Robert *et al.*, 2003) [18]. But this time the invasive *An. Stephensi* has been established in Africa region since 2012. It was first detected in Djibouti in 2012 (Faulde *et al.*, 2014) [12]. Since then, the vector was identified from urban and rural sitting in Ethiopia, Sudan and Somalia in between 2016 and 2019 (Ahmed *et al.*, 2021, WHO 2022 [34], Carter *et al.*, 2018) [3, 34, 10] and further detected from Nigeria (2020), Yemen (2021) and Kenya (2022) (WHO, 2023) [31]. Based on the experience observed from Djibouti, *P. falciparum* incidence might be increased by 50% in Ethiopia due to the invasive *An. Stephensi* (Tadesse *et al.*, 2021, Hamlet *et al.*, 2022) [27, 2]. If the invasion of *anopheles stephesi* continue across the African continent, additional 126 million people from urban areas will be at risk of malaria infection (Sinka *et al.*, 2020) [25]. Recently, (*Ae*) *stephensi* as a competent vector of both *Plasmodium* (*p*). *vivax* and *P. falciparum* appear to be better adapted in urban and peri-urban areas of East African countries (Al-eryani *et al.*, 2023) [6]. It primarily breeds in domestic and peri-domestic household (HH) water containers including water tanks, tyres, basins either plastic or cement, construction sites, barrels and other human-made habitats (Singh *et al.*, 2021, Thomas *et al.*, 2016) [22, 28]. *An. Stephensi* is the primary vector of malaria in Asia (Sinka *et al.* 2011) [23]. Several strategies have been employed in countries endemic with *An. Stephensi*, predominantly targeting the aquatic stages due to the confirmed resistance of this vector to the four major insecticide classes: organochlorines, pyrethroids, organophosphates, and carbamates (Enayati, *et al.* 2020) [11]. In India, Larva control through chemicals and biological means (larvivorous fish and bacteria) are commonly used. Temephos and *Bacillus thuringiensis var israelensis* (*Bti*) are the widely used larvicides (Subbarao, *et al.*, 2019) [26]. According to Al-Eryani *et al.* 2023 [6], the emergence and spread of *An. Stephensi* threatens the progress in malaria control in the Horn of Africa and SSA.

Gash Barka (GB) Region is one of the six Regions in Eritrea. 95% population of the region are living in malaria area. This region is responsible for more than 80% of the total cases produced in Eritrea. According to the annual report from the Communicable Diseases Control (CDC): Eritrea, there was a 57% increase in malaria cases observed between 2015 and 2019, paradoxically contradicting the goals of the 2020-Enitiative: a 38% and 40% reduction in incidence and mortality, respectively. GB records high transmission with annual parasite index of 57.7 in contrast to the other Regions 3.6 in Maekel, 3.8 in Anseba, 9.7 in Debub, 13.8 in southern red sea (SRS) and 7.4 in Northern red sea regions (NRS).

GB Region has shared a border with Ethiopia and Sudan where *An. Stephensi* was incriminated in 2016 and 2019, respectively. The current surveillance effort aims to determine the presence of *An. Stephensi* in the GB Region, using both morphological and molecular identification.

## Methodology

### Study design

Cross sectional entomological survey during the period Sep - Dec. 2022.

**Study area:** Gash-Barka is one of the most malaria zones in the country. It has an altitude range of 142 to 2,702 meters above sea level; it is divided into sixteen administrative sub-zones; and it has mainly a July-September rain season with a small rain season in April/May. The zone is one of the malaria zones in the country. It has a total population of 910, 082, of which 95% (864, 578) live in malaria-endemic areas. Based on recent 2022 malaria data, *Plasmodium falciparum* is the dominant parasite species in the zone, accounting for 88% of infections. *Plasmodium vivax* follows with 12%, and the remaining percentage comprises mixed infections involving either Pf, Pv, or other species. The study was conducted in three towns of the Region: Teseney, Omhajer and Agordet.





Fig 1: Trend of confirmed malaria cases in Tesseney, aqordat and Omhajer from 2015-2022



**Fig 2:** Map of Eritrea, Gash Barka Region and study towns (Aqurdet, Teseney and Golj)

### Tesseney

Tesseney, located in the western part of the Gash Barka Region (GBR), is known as a land port due to its role as a center of trade between Eritrea and its neighboring countries, Sudan and Ethiopia. Tesseney has a total population of 30,000, and its economy relies heavily on farming and animal husbandry. The Gash River runs alongside the town, increasing its vulnerability to malaria transmission. Tesseney is located at  $14^{\circ}19'23''$  N,  $36^{\circ}54'43''$  E, at an elevation of 600 meters above sea level (masl). A dramatic increment in malaria cases was observed between 2015-2022. The number of malaria cases in Tesseney increased more than fivefold, jumping from 727 in 2015 to 3,702 in 2022.

### Agordat

Agordat is the third largest town in GBR, with a population of 22,048. This town has maximum population movement for trade and other economic matters. The economy is heavily driven by agriculture and animal raising, even though malaria cases are increasing. The Gash Barka River, flowing through the town, provides permanent breeding grounds for mosquitoes, contributing to the persistent malaria epidemic despite extensive efforts against both adult and immature mosquitoes. An outstanding achievement was observed in malaria reduction in Eritrea between 2000-2015, but paradoxically a significant increment has been observed since 2015 for unknown reasons. In this town, malaria cases were increased by 624% between 2015-2020 (From 272 to 1699). Agordat located at  $15.551174^{\circ}$  N and  $37.886464^{\circ}$  E at an elevation of 615 masl.

### Omhajer

Omhajer is indeed the closest town to both Ethiopia and Sudan with a population of near 10,000. Omhajer's location at the confluence of borders with Sudan and Ethiopia, combined with unrestricted movement across them, makes it susceptible to cross-border epidemics. Additionally, its proximity to the Tekeze River further increases this

vulnerability. A dramatic change was observed in malaria cases between 2015-2019. It increased by nearly eight fold from 255 to 1985, despite a slight fall in malaria cases was shown in 2022. Omhajer located at  $15.84569^{\circ}$  N,  $37.247269^{\circ}$  E at an elevation of 603.

### Study Population

Larva of *Anopheles* mosquito.

### Sampling Technique and sampling size determination

To identify *Anopheles* mosquito breeding in households (HH), we inspected wet water-holding containers for *Anopheles* larvae and pupae. Collected specimens were then reared to adulthood for morphological and molecular identification.

Any water-holding containers were inspected for the presence of immature mosquitoes (Larva or pupa). To maximize the detection of *Anopheles* mosquito breeding, we inspected all readily accessible water-holding containers within local households (HH). These included basins (Cement, plastic, metal), discarded tins and tires, clay pots, barrels, handwashing basins, leaky faucets, gutters, and other potential breeding sites. A sufficient number of *Anopheles* mosquito immature stages were collected and reared at the malaria entomology laboratory under controlled, optimal environmental conditions.

### Data Collection

During the present survey only immature mosquitoes were used. Larvae and pupa from indoor and outdoor containers were collected either by pipetting or dipping methods. Dipping was done in accordance with the larva searching strategies recommended by WHO (WHO, 1975) [30]. Pipetting was used in small containers not suitable for dipping. The water was first emptied into an enamel tray and local materials and all the larvae and pupae were collected in vials. The collected larva and pupa were brought to the mosquito rearing facility at Tesseney town, where they were placed in trays and raised to adult. Each

enamel tray containing larvae or pupae was labeled by habitat type. The larvae were allowed to develop in the water that was drawn from the field, to maintain the same aquatic environment. The pupae were sorted and transferred with pipettes from the enamel trays to beakers with modest volumes of water, then kept inside cages. A dissecting microscope was used to identify emerging adults to the species using identification key. All identified specimens were preserved individually in Eppendorf tubes for further analysis and were sent to the national malaria control program for further molecular identification.

### Data Analysis

CSPPro7.2 and statistical package for social science (SPSS) software version 21 were used as data entry and analysis respectively. Descriptive statistics were calculated for both variables, and a chi-square test was performed to assess the significance of the association between them. A p-value of less than 0.05 was considered to indicate a statistically significant association. Estimates were given in percentages, odds ratio and confidence intervals.

### Ethical Clearance

Ethical clearance was first obtained from the ethical review committee at the Eritrean Ministry of Health. Upon entering the selected study area, permission was obtained from the sub-zonal administration, municipalities, and local administrations. Additionally, verbal consent was obtained from all household heads.

### Result

#### Malaria morbidity

During the last eight years 28,315 malaria cases were reported from both study areas. *Plasmodium falciparum* (pf) accounts for more than 80% of the cases and the rest were *Plasmodium vivax* (pv) and mixed infection with both pf and pv. Over the past eight years, malaria cases in GB increased nearly fivefold, from 1,224 in 2015 to 5,806 in 2022. (Fig 1).

**Habitat types:** Their positivity and productivity for anopheles and other mosquito type during a 4-month study

period with 8 collection rounds, a total of 828 containers were inspected from three urban sites in Gash Barka zone (Tesseney, Aqordet, and Omhajer). Of these, 186 (22%) were positive for at least one of the three mosquito types. And more than 94% of the total positive containers were found harboring anopheles species (Table 1). The positivity of containers and the number of larva per container were almost the same at all the study sites. Table 3 shows that cement basins and barrels (plastic or metal) were the most common breeding sites for mosquitoes, accounting for over 62% of positive traditional water containers inspected during the study. Although widely used at all study sites, clay pots accounted for only 3% of the total positive mosquito breeding containers.

Table 2: During the study, 18,607 larvae and 1,225 pupae were collected from various container types, including cement basins, plastic basins, 200-liter barrels (plastic and metal), flower vases, clay pots, and overhead tanks. Of these, 8,212 were Anopheles mosquitoes, 7,046 Culex mosquitoes, and 4,349 Aedes mosquitoes. Cement basins stood out as the most productive breeding site, contributing a staggering 87% of collected Anopheles mosquitoes, followed by 45% of Culex and 50% of Aedes species. 200-liter plastic or metal barrels emerged as the second-most productive breeding site, contributing 12% of Anopheles, 31% of Culex, and a notable 42% of Aedes mosquitoes collected during the study. Table 4 reveals that all three mosquito species, Anopheles, Aedes, and Culex, were found to co-breed in the same habitats.

Prevalence of anopheles stephensi

Figure 3 reveals that 383, 242, and 185 female Anopheles mosquitoes were reared for identification from Omhajer, Tesseney, and Agordat, respectively. Each adult mosquito emerged from the pupa was identified to species level using morphological characters under a dissecting microscope two to four days of post-emergence. During morphological identification: 368 (96.1%) from Omhajer, 174 (71.9%) from Tesseney and 105 (56.8%) from Aqordet were found to be *An. Stephensi*. All the distinct features of *An. Stephensi* was observed.

The present result shows the invasive malaria vector known as *An. Stephensi* is well-established at all the study sites.


**Table 1:** Habitat types and their positivity to anopheles larvae in selected sites of the Gash Barka region

Container type	No. of inspected n (%)	Positive for any mosquito larva	Positive for anopheles larva n (%)
Cement basin	178	116 (65%)	109 (94%)
Plastic barrel	312	31 (10%)	16 (52%)
Metal Barrel	143	24 (17%)	10 (42%)
Flower vase	84	2 (2%)	2 (100%)
Clay pot	63	5 (8%)	1 (20%)
Metal tank	45	7 (16%)	5 (55%)
Shower vase	3	1 (33%)	0 (0%)
Total	828	186 (22%)	143 (77%)

**Table 2:** Larvae of anopheles and culicines collected from various habitat type in

Container type	No. of positive containers	No. of containers positive for both mosquito type	No. of anopheles collected	No. of culex collected	No. of aedes collected	No. of pupa
Cement basin	116	38 (31%)	7136	3166	2165	681
Plastic barrel	31	12 (26%)	444	1408	815	380
Metal Barrel	24	5 (21%)	546	800	999	151
Flower vase	2	0	5	200	50	1
Clay pot	5	0	7	195	30	0
Metal tank	7	1(14%)	74	500	240	4
Shower vase	1	0	0	578	50	8
Total	186	56	8212	7046	4349	1225





**Fig 3:** Comparison of anopheles mosquito larvae collected, reared to adult and identified as *An. Stephensi*

## Discussion

The result of the present study presented the morphological identification of *An. Stephensi* in the urban sites of western part of Eritrea. This Region shared a border with Ethiopia and Sudan. *An. Stephensi* was previously identified in almost all the neighboring countries, in Djibouti in 2012 (Faulde *et al.*, 2014) [12], Ethiopia in 2016 (Carter *et al.*, 2018) [1], and later in western Ethiopia in 2022 (Hawaria *et al.*, 2023) [15], in Central Sudan, 2019 (Ahmed *et al.*, 2021) [3], in Somalia (Somaliland) 2019 (Ali *et al.*, 2022) [27], Kenya (Ochomo *et al.*, 2023) [17] and in the in the Middle East like in Yemen (Allan *et al.*, 2023) [8] and Saudi Arabia (Al Ahmad *et al.*, 2011) [5].

The surveillance targets only immature mosquito. No CDC light trap and other adult collection methods were deployed. It was only conducted to ascertain the presence of *An. Stephensi*. Based on the recommendation from WHO, to incorporate *Integrating An. Stephensi* surveillance into the existing malaria vector monitoring system in Eritrea yielded significant results. Surveys conducted in four out of six regions confirmed the presence of this mosquito with a very high prevalence in all urban sites surveyed within Gash Barka region. Molecular identification performed in Sudan by the national malaria control program (NMCP) confirmed the presence of *Anopheles stephensi* in the samples. Unfortunately, detailed information necessary for phylogenetic analysis and comparison with neighboring countries was not provided. Finding phylogenetic similarities with neighboring countries might not be essential in this case, as it's possible that multiple introductions of different clades occurred simultaneously in various parts of Africa. (Ochomo *et al.*, 2023) [17]. The recent surge in malaria cases observed in our study areas over the past five years may be linked to the emergence of this invasive vector. This mosquito species has been demonstrated to be a highly efficient transmitter of both *Plasmodium falciparum* (Pf) and *Plasmodium vivax* (Pv), the parasites responsible for the most severe forms of malaria. A study from Ethiopia suggested that an increment of 50% in malaria incidence could have happened in areas where *An. Stephensi* is prevalent (Hamlet *et al.*, 2022) [2]. A 2020 study by Sinka *et al.* predicted that an additional 126 million people in African urban areas could be exposed to malaria due to the spread of this invasive vector. The 2013 outbreak in Djibouti serves as a stark reminder of how quickly malaria cases can surge. With fewer than 3,000 cases reported in 2013, the number skyrocketed to over

49,000 just five years later (Seyfarth *et al.*, 2019) [19], highlighting the importance of vigilant mosquito control measures.

*An. Stephensi* shares the same breeding habitat with *aedes (Ae.) aegypti*. These mosquitoes prefer HH water holding containers. Cement basins emerged as the dominant breeding site across all study areas, with over 87% of *Anopheles* immatures collected from them. A study in India identified overhead tanks as the dominant breeding site for *Anopheles stephensi*, with over 78% of mosquito larvae collected from this source (Thomas *et al.*, 2016) [22]. All three mosquito species (*Anopheles*, *Culex*, and *Aedes*) were found co-inhabiting the same breeding sites across the study areas, with a relatively even distribution observed. In contrast, a study by Thomas found very limited co-inhabitation of *Anopheles* with other mosquito types in overhead tankers (Less than 5%). This contrasts with our findings of widespread co-inhabitation across different breeding sites.

Unplanned urbanization is creating mosquito-friendly infrastructure, leading to a surge in threats like arboviruses and, more recently, malaria transmitted by *An. Stephensi*, posing significant risks to urban residents. These factors, coupled with climate change creating favorable breeding conditions, pose a significant threat for the spread and establishment of *An. Stephensi* across Africa. (Ochomo *et al.*, 2023) [17]. Furthermore it was believed that this vector is resistant to the four classes of insecticides. In Ethiopia, resistance testing revealed that *Anopheles stephensi* mosquitoes are highly resistant to pyrethroids, carbamates, and organophosphates, which are the main classes of insecticides used for mosquito control (Balkew *et al.*, 2020) [19].

Our understanding of *Anopheles stephensi*'s resting and biting behavior remains limited, despite its growing importance as a malaria vector in Africa. This knowledge gap, as highlighted by Mnzava, hinders effective control strategies (Mnzava *et al.*, 2022) [36]. The finding that *An. Stephensi* bites outdoors challenges the effectiveness of traditional control methods like insecticide-treated nets (ITNs) and indoor residual spraying (IRS), which primarily target indoor-resting mosquitoes (Ochomo *et al.*, 2023) [17]. A community-based intervention targeting the *Aedes aegypti* mosquito, the primary vector of dengue fever, was implemented in Anseba Region, Eritrea. The effectiveness of the intervention, which involved door-to-door education and community mobilization for mosquito breeding site

elimination, was assessed through a reduction in both mosquito populations and reported dengue cases. The community-based approach achieved an impressive 10-fold reduction in larval indices, suggesting its potential as a valuable tool for controlling dengue fever transmission in similar settings (Yenus *et al.*, 2021) [13]. This intervention would be a choice of intervention against *An. Stephensi*, as this vector has the same breeding behavior with *Ae. aegypti*. In addition to this, a social and behavioral change guidance for *An. Stephensi* in Africa was developed and disseminated, which targets to promote the utilization of core interventions (ITS, IRS and prompt care-seeking for fever) and larval source management (Household larviciding, community larviciding, regulatory finding and removing standing water and covering water storage container).

### Conclusion and Recommendation

*Anopheles stephensi* co-occurred with various native vector species, including [culex and aedes], at all three study sites. Further research is needed to determine its relative role in malaria transmission in the region. Given the co-existence of *Anopheles stephensi* with native malaria vectors, health authorities urgently need to revise existing control strategies. This comprehensive approach must simultaneously target both dominant and newly identified vectors to effectively manage their behavior and prevent disease transmission.

### References

1. Carter TE, Yared S, Gebresilassie A, Bonnell V, Damodaran L, Lopez K, *et al.* First detection of *Anopheles stephensi* Liston, 1901 (Diptera: Culicidae) in Ethiopia using molecular and morphological approaches. *Acta Trop.* 2018;188:180-186.
2. Hamlet A, Dengela D, Tongren JE, Tadesse FG, Bousema T, Sinka M, *et al.* The potential impact of *Anopheles stephensi* establishment on the transmission of *Plasmodium falciparum* in Ethiopia and prospective control measures. *BMC Med.* 2022;20:135.
3. Ahmed A, Khogali R, Elnour MB, Nakao R, Salim B. Emergence of the invasive malaria vector *Anopheles stephensi* in Khartoum State, Central Sudan. *Parasit Vectors.* 2021;14:511.
4. Ahmed A, Khogali R, Elnour MAB, Nakao R, Salim B. Emergence of the invasive malaria vector *Anopheles stephensi* in Khartoum State, Central Sudan. *Parasites & Vectors.* 2021;14(1):1-5.
5. Al Ahmad A, Sallam M, Khuriji M, Kheir S, Azari-Hamidian S. Checklist and pictorial key to fourth-instar larvae of mosquitoes (Diptera: Culicidae) of Saudi Arabia. *J Med Entomol.* 2011;4:17-37.
6. Al-Eryani SM, Irish SR, Carter TE, Lenhart A, Aljasari A, Montoya LF, *et al.* Public health impact of the spread of *Anopheles stephensi* in the WHO Eastern Mediterranean Region countries in Horn of Africa and Yemen: need for integrated vector surveillance and control. *Malaria Journal.* 2023;22(1):187.
7. Ali S, Samake JN, Spear J, Carter TE. Morphological identification and genetic characterization of *Anopheles stephensi* in Somaliland. *Parasites & Vectors.* 2022;15(1):247.
8. Allan R, Weetman D, Sauskojus H, Budge S, Hawaii TB, Baheshm Y. Confirmation of the presence of *Anopheles stephensi* among internally displaced people's camps and host communities in Aden city, Yemen. *Malaria Journal.* 2023;22(1):1-8.
9. Balkew M, Mumba P, Dengela D, Yohannes G, Getachew D, Yared S, *et al.* Geographical distribution of *Anopheles stephensi* in eastern Ethiopia. *Parasit Vectors.* 2020;13:35.
10. Carter TE, Yared S, Gebresilassie A, Bonnell V, Damodaran L, Lopez K, *et al.* First detection of *Anopheles stephensi* Liston, 1901 (Diptera: Culicidae) in Ethiopia using molecular and morphological approaches. *Acta Trop.* 2018;188:180-186.
11. Enayati A, Hanafi-Bojd AA, Sedaghat MM, Zaim M, Hemingway J. Evolution of insecticide resistance and its mechanisms in *Anopheles stephensi* in the WHO Eastern Mediterranean Region. *Malar J.* 2020;19:258.
12. Faulde MK, Rueda LM, Khaireh BA. First record of the Asian malaria vector *Anopheles stephensi* and its possible role in the resurgence of malaria in Djibouti, Horn of Africa. *Acta Trop.* 2014;139:39-43.
13. Mensur Y, Micheal B, Mickeal B, Amanuel K, Kiros S. A comprehensive analysis on abundance, distribution and bionomics of potential malaria vectors in Anseba zone, Eritrea. *Int. J Mosq Res.* 2021;8(2):116-124.
14. Hamlet A, Dengela D, Tongren JE, Tadesse FG, Bousema T, Sinka M, *et al.* The potential impact of *Anopheles stephensi* establishment on the transmission of *Plasmodium falciparum* in Ethiopia and prospective control measures. *BMC Med.* 2022;20:135.
15. Hawaria D, Kibret S, Zhong D, Lee MC, Lelisa K, Bekele B, *et al.* First report of *Anopheles stephensi* in Southern Ethiopia.
16. Himeidan YE, Rayah EE. Role of some environmental factors on the breeding activity of *Anopheles arabiensis* in New Halfa town, eastern Sudan. *East Mediterr Health J.* 2008;14:252-9.
17. Ochomo EO, Milanoi S, Abong'o B, Onyango B, Muchoki M, Omoke D, *et al.* Molecular surveillance leads to the first detection of *Anopheles stephensi* in Kenya.
18. Robert V, *et al.* Malaria transmission in urban sub-Saharan Africa. *Am J Trop Med Hyg.* 2003;68:169-176.
19. Seyfarth M, Khaireh BA, Abdi AA, Bouh SM, Faulde MK. Five years following first detection of *Anopheles stephensi* (Diptera: Culicidae) in Djibouti, Horn of Africa: populations established-malaria emerging. *Parasitol Res.* 2019;118:725-732.
20. Shilulu J, Mbogo C, Ghebremeskel T, Githure J, Novak R. Mosquito larval habitats in semiarid ecosystems in Eritrea: impact of larval habitat management on *Anopheles arabiensis* population. *Am J Trop Med Hyg.* 2007;76:103-10.
21. Shilulu J, Ghebremeskel T, Mengistu S, Fekadu H, Zerom M, Mbogo C, *et al.* Distribution of *Anopheles* mosquitoes in Eritrea. *Am J Trop Med Hyg.* 2003;69(3):295-302.
22. Singh H, Gupta SK, Vikram K, Saxena R, Sharma A. The impact of mosquito proof lids of underground tanks tanka on the breeding of *Anopheles stephensi* in a village in western Rajasthan. *India Malar J.* 2021;20:412.
23. Sinka ME, Bangs MJ, Manguin S, Chareonviriyaphap T, Patil AP, Temperley WH, *et al.* The dominant *Anopheles* vectors of human malaria in the Asia-Pacific

- region: occurrence data, distribution maps and bionomic précis. *Parasit Vectors*. 2011;4:89.
24. Sinka ME, Bangs MJ, Manguin S, Coetzee M, Mbogo CM, Hemingway J, *et al.* The dominant *Anopheles* vectors of human malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and bionomic précis. *Parasit Vectors*. 2010;3:117.
  25. Sinka ME, Pironon S, Massey NC, Longbottom J, Hemingway J, Moyes CL, *et al.* A new malaria vector in Africa: Predicting the expansion range of *Anopheles stephensi* and identifying the urban populations at risk. Check for updates, University of Oslo, Norway. 2020;177/40.
  26. Subbarao SK, Nanda N, Rahi M, Raghavendra K. Biology and bionomics of malaria vectors in India: existing information and what more needs to be known for strategizing elimination of malaria. *Malar J*. 2019;18:396.
  27. Tadesse FG, Ashine T, Teka H, Esayas E, Messenger LA, Chali W, *et al.* *Anopheles stephensi* mosquitoes as vectors of *Plasmodium vivax* and *Plasmodium falciparum*, Horn of Africa, 2019. *Emerg Infect Dis*. 2021;27:603-607.
  28. Thomas S, Ravishankaran S, Justin JA, Asokan A, Mathai MT, Valecha N, *et al.* Overhead tank is the potential breeding habitat of *Anopheles stephensi* in an urban transmission setting of Chennai, India. *Malar J*. 2016;15:274.
  29. Thomas S, Ravishankaran S, Justin JA, Asokan A, Mathai MT, Valecha N, *et al.* Overhead tank is the potential breeding habitat of *Anopheles stephensi* in an urban transmission setting of Chennai, India. *Malar J*. 2016;15:1-10.
  30. World Health Organization. Manual on practical entomology: part II method and techniques. Geneva: World Health Organization; c1975.
  31. World Health Organization. Malaria Threats Map. Geneva, World Health Organization; c2023. Available from: <https://apps.who.int/malaria/maps/threats>
  32. World Health Organization. World malaria report 2021. Regional data and trends. Goals and milestones of the global technical strategy for malaria 2016-2030.
  33. World Health Organization. Zeroing in on malaria elimination; c2021: final report of the E-2020 initiative.
  34. World Health Organization. World malaria report 2022. Geneva, World Health Organization; c2022. Available from: <https://apps.who.int/iris/handle/10665/365169>
  35. World Health Organization. WHO initiative to stop the spread of *Anopheles stephensi* in Africa. Geneva, Switzerland: World Health Organization; c2022.
  36. Mnzava A, Monroe AC, Okumu F. *Anopheles stephensi* in Africa requires a more integrated response. *Malaria journal*. 2022 May 31;21(1):156.